

## CHEMICAL POLLUTANTS IMPACT IN PLANTS: PHYSIOLOGICAL AND MOLECULAR STUDY - A REVIEW

Basel Saleh\*

*Department of Molecular Biology and Biotechnology, AECS, P.O. Box 6091, Damascus, Syria*  
*Basel Saleh, PhD. Department of Molecular Biology and Biotechnology, Atomic Energy Commission of Syria, P.O. Box 6091, Damascus, Syria.*

(Received on Date: 24<sup>th</sup> January 2017

Date of Acceptance: 6<sup>th</sup> March 2017)

### ABSTRACT

Huge experiments on pollution monitoring have been realized to evaluate pollution impact on marine and terrestrial environments using physiological, biochemical and molecular aspects. Thereby, many plants species were used as a potent bioindicator for monitoring pollution with low cost. Employment of algae as bioindicator for detection environmental pollution in marine ecosystems has been attained recently much attention. Chemical pollutants impacts on living organism (bacteria, plants and animals) could manifest by growth impair, pigments content reduction, organism death and DNA damages and mutations. Their dangers occurred in terrestrial and aquatic environmental ecosystems, in different ways. The current investigation mainly reported chemical pollutants impacts on plants due to their highly potent in accumulation these pollutants in their tissues.

**Keywords:** Chemical pollutants, Marine Algae, Plants, Toxicity, Genotoxicity

---

**No of Tables : 2**

**No of References : 77**

---

## INTRODUCTION

Heavy metal contamination from the dangers that threaten the environmental ecosystems including seas, lakes and rivers worldwide. The interest of pollutants worldwide increased; these being of high toxicity on one hand and does not break down naturally in the environment on the other hands. Therefore, many studies have focused on pollutants behavior investigation, their toxicity and ability to move on to the food chain and possible ways to remove it. This phenomenon may pose a direct threat to humans and public health (Phillips and Rainbow, 1993; Rainbow, 1995; Bing *et al.*, 2013; Khan *et al.*, 2013).

Pollution in marine ecosystems is a serious environmental issue. It considerably increased due to augmentation human industrial activities. Thereby, monitoring of chemical pollutants in marine ecosystems became one of the indispensable interests in environment. Even, use of living organisms for detection marine pollution could be considered as a potent tool and provide early warning signals of water pollution trends with low cost. Heavy metals impact has been widely investigated in board reports. In this regards, Jamers *et al.* (2013) reported that, cadmium (Cd), mercury (Hg) and lead (Pb) are considered by the US Environmental Protection Agency (EPA) the three contaminants of greatest threat to the environment. Their impacts on living organisms have been widely investigated at physiological, biochemical and

molecular levels. Many reports showed their toxicity in terrestrial (Shahrtash *et al.*, 2010; Liu *et al.*, 2012; Azimi *et al.*, 2013; Zhang *et al.*, 2015; Salarizadeh and Reza-Kavousi, 2015), aquatic (Gupta and Sarin, 2009) and marine (Wang *et al.*, 1998; Atienzar *et al.*, 2000; Zhou *et al.*, 2011; Bouzon *et al.*, 2012; Saleh, 2015a; 2016b; 2016e) environmental ecosystems.

Previously, it has been reported that heavy metal stress caused DNA changes manifested by loss or induced new bands compared to their respective control. Thereby, DNA alteration patterns could be manifested by disappearance of control bands (DNA damage) or induction of new bands (DNA mutation) (Labra *et al.*, 2003; Atienzar and Jha, 2006; Saleh, 2015b, 2016a; 2016c & 2016d). In this respect, Labra *et al.* (2003) stated that RAPD or AFLP techniques were most potent than classic genotoxic assays, even RAPD marker could serve as an efficient tool for detecting temporary DNA changes that might not consequently manifest themselves as mutations. It has been demonstrated that PARD assay among of molecular markers available, could be considered as a potent and novel biomarker for monitoring DNA changes even DNA mutations induced by adverse environmental stresses in plants, animals, bacteria and algae (Wong *et al.*, 2000; Liu *et al.*, 2005; 2007; Aksakal *et al.*, 2013; Aksakal and Esim, 2015; Saleh, 2015b, 2016a, 2016c; 2016d).

Since 2014, Hori *et al.* reported that the terrestrial *Klebsormidium flaccidum* green algae shares some genes involved in light and drought tolerance with terrestrial plants; based on cell wall traits combined with the recently sequenced genome.

Thereby, the current review mainly focused on physiological and genetic DNA changes induced by heavy metals stress in plants and algae using molecular marker tools. Consequently, it will be somewhat gave an overview image about the importance of some living organisms (plants and algae) as a potent bioindicator for monitoring pollution in ecosystems.

### Potential algae use

Algae displayed an multiuse role in different manner; e.g. as bioindicator for heavy metal pollution (Rainbow, 1995); heavy metals biosorption (Vilar *et al.*, 2008); bioremediation of wastewater

(Prabha *et al.*, 2016); as antimicrobial agent (Saleh and Al-Mariri, 2016).

### Physiological approach

To date, toxicity of different chemical pollutants on living organisms at physiological level have been successfully documented in several researches (Table 1).

Cadmium (Cd) among chemical pollutants displayed an important role as a potent toxicant to bacteria, algae, and fungi, where algae were the most sensitive (Trevors *et al.*, 1986; Saleh, 2016c).

**Table 1.** Chemical pollutants toxicity in living organisms at physiological approach.

Living organisms	Metal ion	Reference
<b>Plants species</b>		
<b>Monocotyledons</b>		
Wheat ( <i>Triticum aestivum</i> L.)	Cd	Milone <i>et al.</i> (2003)
Wheat ( <i>Triticum aestivum</i> L.)	B	Kekec <i>et al.</i> (2010)
Wheat ( <i>Triticum aestivum</i> L.)	Pb	Pazoki <i>et al.</i> (2014)
		Verma and Dubey (2001)
Rice ( <i>Oryza sativa</i> L.)	Cd	Shah <i>et al.</i> (2001)
Rice ( <i>Oryza sativa</i> L.)	Cd	Maksimovic <i>et al.</i> (2007)
Maize ( <i>Zea mays</i> L.)	Cd and Ni	Erturk <i>et al.</i> (2015)
Maize ( <i>Zea mays</i> L.)	B and Zn	Gupta and Sarin (2009)
<i>Hydrilla verticillata</i>	Cd, Hg and Cu	

**Dicotyledons**

<i>Iris pseudacours</i> L.	Cr and Zn	Caldelas <i>et al.</i> (2012) Jana and Choudhuri (1984)
Submerged aquatic plants	Hg, Pb, Cd and Cu	(1984)
Okra ( <i>Abelmoschus esculantus</i> L.)	Cd	Aydin <i>et al.</i> (2013)
Chili peppers ( <i>Capsicum annum</i> L.)	Cd	Aslam <i>et al.</i> (2014) Ortega-Villasante <i>et al.</i> (2005)
Alfalfa ( <i>Medicago sativa</i> L.)	Cd and Hg	Gichner <i>et al.</i> (2004)
Tobacco ( <i>Nicotiana tabacum</i> L.)	Cd	
<i>Sesuvium portulacastrum</i> and <i>Mesembryanthemum crystallinum</i>	Cd	Ghnaya <i>et al.</i> (2007)
Pea ( <i>Pisum sativum</i> L.)	Cd	Fusconi <i>et al.</i> (2007)
<i>Kalanchoe</i>	Cd	Ozyigit <i>et al.</i> (2016a)
Bean ( <i>Phaseolus vulgaris</i> L.)	Ni, Co, Cr and Zn	Zengin (2013) Gupta and Sarin (2009)
<i>Ceratophyllum demersum</i>	Cd, Hg and Cu	
<b>Algae</b>		
<i>Chlorella</i> sp (Chlorophyta)	Cd	Kaplan <i>et al.</i> (1995) Amado Filho <i>et al.</i> (1996)
<i>Padina gymnospora</i> (Phaeophyta)	Cu and Zn	Amado Filho <i>et al.</i> (1997)
Six examined seaweeds (C, R and Ph)	Zn	Lewis <i>et al.</i> (1998)
<i>Enteromorpha intestinalis</i> (Chlorophyta)	Cu	Malea <i>et al.</i> (2006)
<i>Enteromorpha</i> spp. (Chlorophyta)	Cd, Zn and N	
<i>Potamocorbula amurensis</i> and <i>Macoma balthica</i>	Cd, Cr, and Zn	Lee <i>et al.</i> (1998)
<i>Ulva lactuca</i> (Chlorophyta)	Cr(VI)	Unal <i>et al.</i> (2010)
<i>Ulva lactuca</i> (Chlorophyta)	Cu, Pb, Cd and Zn	Saleh (2015a)
<i>U. lactuca</i> (Chlorophyta)	Pb	Saleh (2016e)
<i>U. lactuca</i> (Chlorophyta) and <i>Padina pavonica</i> (Phaeophyta)	Cd	Saleh (2016b)
<i>Ulva lactuca</i> (Chlorophyta)	Cd	Markham <i>et al.</i> (1980) Bulgariu and Bulgariu (2012).
<i>Ulva lactuca</i> (Chlorophyta)	Pb, Cd and Co	Jiang <i>et al.</i> (2013)
<i>U. prolifera</i> and <i>U. linza</i> (Chlorophyta)	Cd	Vilar <i>et al.</i> (2008)
<i>Gelidium sesquipedale</i> (Rhodophyta)	Cu, Pb, Cd and Zn	Collén <i>et al.</i> (2003)
<i>Gracilaria tenuistipitata</i> (Rhodophyta)	Cu	Xia <i>et al.</i> (2004)
<i>Gracilaria lemaneiformis</i> (Rhodophyta)	Cu and Cd	dos Santos <i>et al.</i> (2012)
<i>Gracilaria domingensis</i> (Rhodophyta)	Cd	

<i>Hypnea musciformis</i> (Rhodophyta)	Cd	Bouzon <i>et al.</i> (2012)
<i>Nitzschia closterium</i> (Bacillariaceae)	Cd, Ni, Pb, Zn, Fe and CrVI	Ova and Övez (2013)

It has been demonstrated that, some elements e.g. copper (Cu), zinc (Zn) and iron (Fe) in low concentrations are essential for catalyzing enzymatic reactions in living organisms. Other elements however, e.g. lead (Pb), mercury (Hg) and cadmium (Cd) could be associated as cofactors for activation of their enzymatic systems (Manoj and Padhy, 2013; Saleh, 2015a; 2016b; 2016e). Chemical pollutants can affect the plant and algal growth and physiological and biochemical activities; e.g. Photosynthesis activity and chlorophyll content. These effects could be manifested by decline in specific growth rate SGR%, pigmentations content (chlorophyll and carotenoids) and total protein content (Bouzon *et al.*, 2012; dos Santos *et al.*, 2012; Jiang *et al.*, 2013).

It has been reported that Cd and Pb toxicity in chlorophyll content in plants could be attributed to their role in inhibition of key enzymes such as d-aminolevulinic acid dehydratase (ALA-dehydratase) and protochlorophyllide reductase associated with chlorophyll biosynthesis (Aslam *et al.*, 2014). Indeed these metals could impair the supply with Mg, Fe and Zn (Kupper *et al.*, 1996; Aslam *et al.*, 2014). More recently, Ozyigit *et al.* (2016b) investigated Pb (0, 100, 200 and 400  $\mu\text{mol/L}$ ) impact in rye (*Secale cereale* L.) after 2 weeks exposure. The previous study showed decline in physiological parameters e.g. chlorophyll a, chlorophyll b, total chlorophyll and

carotenoids content by 6.68%, 6.08%, 2.89% and 8.57%, respectively at the highest Pb applied concentration (400  $\mu\text{mol/L}$ ). Indeed, Erturk *et al.* (2015) reported boron (B) and zinc (Zn) impact on total soluble protein content in *Z. mays*. The previous study revealed that Zn ion reduced the previous parameter and this decline was more noticeable with Zn than B ions.

Moreover, Gupta and Sarin (2009) reported 10  $\mu\text{mol/L}$  Cd, 5  $\mu\text{mol/L}$  Hg, and 20  $\mu\text{mol/L}$  Cu impacts on *H. verticillata* and *C. demersum* aquatic plants for 96 h. The previous study revealed that heavy metal applied reduced chlorophyll and protein contents into the two species. This reduction was more pronounced in *H. verticillata* compared to *C. demersum*. Whereas, Caldelas *et al.* (2012) reported physiological response of *Iris pseudacorus* L. plants to chromium ( $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$ ) and zinc ( $\text{ZnCl}_2$ ) (0, 10, 50, 100, and 200  $\mu\text{g/mL}$ ) for 5 weeks. The previous study revealed growth impair by 65% and 31% (dry weight) with Cr and Zn treatment, respectively. Whereas, metal treatment has no effect on photosynthetic pigment content. While, Aslam *et al.* (2014) investigated Cd (20, 40, 60, 80 and 100 ppm) impact on *C. annuum* L. for 90 days exposure. The previous study showed that Cd stress caused significant reduction in shoot length, root length and fresh weight, protein, chlorophyll and carotenoid content.

More recently, Ozyigit *et al.* (2016a) reported that toxicity of Cd in *Kalanchoe* plants *in vitro* for two months. The previous study revealed that 400  $\mu$ M Cd caused reduction in Cha, Chb, Total Chl and carotenoids pigments recorded by ~40.57, ~37.63, ~36.27 and ~37.66%, respectively compared to their respective control.

As for algae, Unal *et al.* (2010) reported response of *U. lactuca* to chromium (VI) (0.2, 0.5, 1, and 5 mM  $K_2CrO_4$ ) after 2 h Cr(VI) exposure. The previous study revealed morphological changes in thallus cells, decline in photochemical efficiency of PSII (Fv/Fm) ratio combined with increase in necrotic cells along with increased of Cr(VI) from 1 to 5 mM Cr(VI), than their respective control. Whereas, Jiang *et al.* (2013) reported none significant decline in both chlorophyll and carotenoid pigments observed under 1.8 mg/L Cd compared to their respective control in both *U. prolifera* and *U. linza* marine algae. While, with 3.7, 7.4 and 14.8 mg/L Cd, a decline in Chl content by 18, 25 and 45% in *U. prolifera*, respectively has been observed; and it was 16, 20 and 39% for *U. linza*, respectively. As for carotenoids, this decline was found to be 16, 29 and 54%, respectively in the case of *U. prolifera* and by 13, 16 and 44% below the control for *U. linza*, respectively.

Whereas, Saleh (2015a) reported Cu (5.8 mg/L), Pb (18.2 mg/L), Cd (10.5 mg/L) and Zn (9.9 mg/L) physiological impacts on *U. lactuca* after 5 days stress. The previous study showed that metal stress reduced SGR% in the following order: Cd > Pb > Zn &

Cu. Whereas, decline in Chla by 66%, 64%, 53% and 50% with Pb, Cd, Cu and Zn treatment, respectively was recorded. While, Pb treatment caused the highest reduction in Chlb reduced by 80% compared to other tested ions. More recently, Saleh (2016e) studied toxicity of Pb (0, 2, 4 and 8 mg/L) after 2 days exposure in *U. lactuca*. The previous investigation revealed that Chla decreased by 1.32, 22.7 and 40.4% at 2, 4 and 8 mg/L Pb. Whereas, this decline was recorded to be 15.8, 13.4 and 17.7% for Chlb below their respective control under the above mentioned Pb concentrations. Moreover, Saleh (2016b) reported Cd (0, 2.5, 5 and 10 mg/L) toxicity in *U. lactuca* (Chlorophyta) green and *P. pavonica* (Phaeophyta) brown seaweeds for 4 days. The previous study could suggest that *U. lactuca* was most sensitive to Cd stress than *P. pavonica*; by showing highest decline in the examined physiological indices than *P. pavonica*. The previous study showed that Cd stress caused decline in Chla ranged from 56.123 to 66.865% in *U. lactuca* and from 16.183 to 21.365% in *P. pavonica* when Cd applied concentration increased from 2.5 to 10 mg/L, respectively. Whereas, carotenoids content decreased from 36.788 to 44.058% in *U. lactuca* when Cd applied concentration increased from 2.5 to 10 mg/L, respectively. While, the previous parameter increased from 0.687 to 80.308% in *P. pavonica* as Cd applied concentration increased from 2.5 to 10 mg/L, respectively. Increased observed carotenoids content in *P. pavonica* as Cd applied concentration increased, makes

them more tolerant to Cd stress compared to *U. lactuca*. Carotenoids pigments act as antioxidant and thereby, serve as osmolytes involved in heavy metal detoxification by minimize ROS induction generated by heavy metals stress (dos Santos *et al.*, 2012; Bouzon *et al.*, 2012).

Similarly, Zengin (2013) reported that carotenoids content significantly increased in the bean (*P. vulgaris* L.) leaves at different concentrations of Ni, Co, Cr and Zn heavy metals. Whereas, Pazoki *et al.* (2014) reported that Pb (0, 300, 600 and 900 mg/kg of soil) reduced chlorophyll *a* and *b* with increases carotene content in wheat (*T. aestivum* L.) as Pb concentration increased from 300 to 900 mg/kg. Previously, Kaplan *et al.* (1995) reported the induction of phytochelatins in *Chlorella* sp. algae subjected to Cd. Moreover, Shariati and Yahyaabadi (2006) reported a decline in chlorophyll content and increase of beta-carotene pigmentations in the two strains (Iranian and Australian) of *Dunaliella salina* green algae as Cd concentration increased from 0.005 to 0.5 mg/L, after 5 days exposure to Cd.

Overall, according to Saleh (2016d), marine algae differently responded to Cd metal than the higher plants. This observation could be related to the absence of root system in algae and its presence in higher plants on one hand. On the other hands, the interaction between

metal and these two living organisms is completely different. The occurrence of algae in direct contact with metal in metal solution make them differently responded against heavy metal stress. Whereas, in terrestrial plants, Cd absorbed in different manner. Consequently, leading algae to adopt different protective mechanisms to minimize or detoxification unfavorable effect of applied metal.

### Molecular approach

Previously, it has been documented that heavy metals pollutants caused oxidative and carcinogenic effects to living organisms in ecosystems. Their toxicity comes from its binding with nucleic acids through various reactions (direct or/and indirect) with DNA sites as well as affecting DNA replication. These phenomena lead consequently to DNA damages and mutations (Valavanidis and Vlachogianni, 2010). These chemical pollutants provoked free radicals and reactive oxygen species (ROS), which arbitrary attack and damage DNA and important enzymatic proteins (Bal and Kasprzak, 2002). Lead (Pb) is one of fewer heavy metals beside Hg and Cd, classified as high toxic element for living organisms in environmental ecosystems (Valavanidis and Vlachogianni, 2010). Many investigations reported DNA changes induced by chemical pollutants in living organisms (Table 2).

**Table 2.** Genotoxicity effect of chemical pollutants in living organisms as revealed by different applied molecular markers.

Living organisms	Metal ion	Molecular marker	Reference
<b>Plants species</b>			
<b>Monocotyledons</b>			
Wheat ( <i>Triticum aestivum</i> L.)	B	RAPD	Kekec <i>et al.</i> (2010)
Wheat ( <i>Triticum aestivum</i> L.)	Cd	RAPD	Azimi <i>et al.</i> (2013)
Barley ( <i>Hordium vulgare</i> L.)	Cd	RAPD	Liu <i>et al.</i> (2005)
Rice ( <i>Oryza sativa</i> L.)	Cd	AFLP	Aina <i>et al.</i> (2007)
Rice ( <i>Oryza sativa</i> L.)	AS (III)	RAPD	Ahmad <i>et al.</i> (2012)
Rice ( <i>Oryza sativa</i> L.)	Cd	SRAP	Zhang <i>et al.</i> (2015)
Maize ( <i>Zea mays</i> L.)	Cd	RAPD	Shahrtash <i>et al.</i> (2010)
Maize ( <i>Zea mays</i> L.)	fungicide	RAPD	Aksoy <i>et al.</i> (2015)
Maize ( <i>Zea mays</i> L.)	sportak	RAPD	Erturk <i>et al.</i> (2015)
Maize ( <i>Zea mays</i> L.)	B and Zn	RAPD	Gupta and Sarin (2009)
Maize ( <i>Zea mays</i> L.)	Cd, Hg and	RAPD	
Maize ( <i>Zea mays</i> L.)	Cu	and SCAR	
<i>Hydrilla verticillata</i>			
<b>Dicotyledons</b>			
Okra ( <i>Abelmoschus esculantus</i> L.)	Cd	RAPD	Aydin <i>et al.</i> (2013)
Cucumber ( <i>Cucumis sativus</i> L.)	Cu and Zn	RAPD	Aydin <i>et al.</i> (2012)
Cucumber ( <i>Cucumis sativus</i> L.)			Salarizadeh and Reza-Kavousi (2015)
Cumin ( <i>Cuminum cyminum</i> L.)	Cd	RAPD	Körpe and Aras (2011)
Eggplant ( <i>Solanum melongena</i> L.)	Cu	RAPD	Aydin <i>et al.</i> (2015)
Tomato ( <i>Lycopersicon esculentum</i> L.)	Pb	RAPD	Ozyigit <i>et al.</i> (2016b)
Rye ( <i>Secale cereal</i> L.)	Pb	RAPD	Ozyigit <i>et al.</i> (2016a)
Kalanchoe	Cd	RAPD	Al-Khateeb and Al-Qwasemeh (2014)
<i>Solanum nigrum</i> L. (wild relative for tomato)	Cd, Cu and Zn	ISSR	Zhan <i>et al.</i> (2011)
<i>Arabidopsis thaliana</i>	Cd	RAPD	Liu <i>et al.</i> (2012)
<i>Arabidopsis thaliana</i>	Cd	RAPD	Ghiani <i>et al.</i> (2014)
White clover	Cd and AS	RAPD	Aly (2012)
Egyptian clover and Sudan grass	Cd	RAPD	Enan (2006)
Common bean ( <i>Phaseolus vulgaris</i> L.)	Cd, Pb, Mn	RAPD	



	Hg, Cr and Zn	RAPD	Cenkci <i>et al.</i> (2009)
Common bean ( <i>Phaseolus vulgaris</i> L.)	Al and Ni	RAPD	Al-Qurainy (2009)
Common bean ( <i>Phaseolus vulgaris</i> L.)	B	RAPD	Kekec <i>et al.</i> (2010)
Rucola ( <i>Eruca sativa</i> L.)	Cd, Pb and Zn	ISSR	Al-Qurainy (2010)
Rucola ( <i>Eruca sativa</i> L.)	Cd, Pb and Zn	RAPD	Al-Qurainy <i>et al.</i> (2010)
Artichoke ( <i>Cynara scolymus</i> L.) and Runner bean ( <i>Phaseolus coccineus</i> L.)	Pb	RAPD	Candan and Batir (2015)
<i>Ceratophyllum demersum</i>	Cd, Hg and Cu	RAPD and SCAR	Gupta and Sarin (2009)
Mung bean ( <i>Vigna radiata</i> L.)	Tannery effluents (CETP)	RAPD	Raj <i>et al.</i> (2014)
Chili peppers ( <i>Capsicum annum</i> L.)	Cd	RAPD	Aslam <i>et al.</i> (2014)
Water Lettuce ( <i>Pistia stratiotes</i> )	Pb	ISSR	Neeratanaphan <i>et al.</i> (2014)
<b>Algae</b>			
<i>Ulva lactuca</i> (Chlorophyta)	Cu, Pb, Cd and Zn	RAMP	Saleh (2015b)
<i>Ulva lactuca</i> (Chlorophyta)	Cu, Pb, Cd and Zn	RAMP	Saleh (2016a)
<i>Ulva lactuca</i> (Chlorophyta)	Cd	RAPD	Saleh (2016c)
<i>Padina pavonica</i> (Phaeophyta)	Cd	RAPD	Saleh (2016d)

Heavy metals genotoxicity as expressed in DNA changes patterns could be manifested by estimated genomic template stability (GTS%) as a qualitative measurement of DNA damage induced by pollutants as reported in many investigations (Liu *et al.*, 2005; Cenkci *et al.*, 2010; Aydin *et al.*, 2012; 2013; Salarizadeh and Reza-Kavousi, 2015; Zhang *et al.*, 2015; Saleh, 2016c; 2016d).

More recently, Ozyigit *et al.* (2016b) reported Pb (0, 100, 200 and 400  $\mu\text{mol/L}$ ) genotoxicity in rye (*S. cereale* L.) after 2 weeks exposure, using RAPD marker. The

previous study revealed RAPD profile alterations and that loss and gains bands were positively correlated with Pb applied concentrations. In this regards, Pb treatment induced 3 newly bands appearance at 100 and 200  $\mu\text{mol/L}$  Pb combined with loss of 2 bands at 200 and 400  $\mu\text{mol/L}$  compared to their respective control. Moreover, GTS% values decreased as Pb applied concentration increased.

Erturk *et al.* (2015) reported boron (B) and zinc (Zn) genotoxicity on *Z. mays* using 16 RAPD primers. The previous investigation

showed that GTS% decreased as Zn and B applied concentrations increased.

Gupta and Sarin (2009) reported 10 mol/L Cd, 5 mol/L Hg, and 20 mol/L Cu impacts on *H. verticillata* and *C. demersum* aquatic plants for 96 h, using RAPD and sequence characterized amplified region (SCAR) markers. The previous investigation showed that heavy metal stress induced DNA profile alteration expressed as reduction in GTS%. This reduction was more noticeable in *H. verticillata* than *C. demersum* one.

Aslam *et al.* (2014) detected DNA changes induced by Cd stress in *C. annuum* L. using 10 RAPD primers. The previous study revealed 184 bands of which 62 bands were polymorphic represent polymorphism level of 34%.

More recently, Saleh (2016a) applied Random amplified microsatellite polymorphism (RAMP) marker for detection of genetic DNA alteration in the green algae *U. lactuca* 5 days after Cu, Pb, Cd and Zn exposure. The previous investigation revealed that GTS% value was recorded to be 65.215, 64.630, 59.835 and 59.250% for Pb, Cu, Cd and Zn ions, respectively.

Moreover, Saleh (2016c) reported Cd-genotoxicity in *U. lactuca* green algae exposed to Cd stress for 4 days using 20 RAPD primers. The previous study showed that GTS% increased from 45.4 to 72.8% as Cd increased from 2.5 to 10 mg/L Cd. Similarly, more recently, Saleh (2016d) reported same metal stress treatment in *P. pavonica* brown marine algae using 22 RAPD primers. The previous investigation revealed that GTS% value increased also from 30.7 to 42.7% as Cd applied

concentration increased from 2.5 to 10 mg/L Cd.

These observations could be suggested that the interaction between RAPD pattern alterations and Cd applied concentrations was dose-dependent. Similarly, Liu *et al.* (2005) reported similar trends in barley (*H. vulgare*) and Aydin *et al.* (2013) in Okra (*A. esculantus* L.) under Cd stress using RAPD marker.

Indeed, Aydin *et al.* (2012) reported similar findings in Cu and Zn genotoxicity of Cucumber (*C. sativus* L.) using RAPD marker. The previous investigation revealed that GTS% increased from 15.98 to 38.69% as ions concentration increased from 40 to 640 mg/L. Moreover, Aydin *et al.* (2013) studied Cd genotoxicity in Okra (*A. esculantus* L.) after 21 Cd exposure, using RAPD marker. The previous study showed that GTS% increased from 59 to 72.5% as Cd applied concentrations increased from 30 to 120 mg/L Cd. Whereas, Salarizadeh and Reza-Kavousi (2015) reported that the GTS% values decreased in *C. cyminum* as Cd applied concentration increased from 300 to 1050  $\mu$ M after 7 days Cd treatment using 10 RAPD primers. These findings could be explained by the fact; decreased GTS% does not related to high DNA damages. Increased GTS% as Cd applied concentrations increased, could be serve as a defense mechanism leading to best and effective DNA repairing. This observation could be related to the fact, that the DNA changes in RAPD profiles appearance of new or/and disappearance of normal control bands);

tendency to balance each other (Liu *et al.*, 2005, Aydin *et al.*, 2013; Saleh, 2016c; 2016d).

Meanwhile, other reports however mentioned an inverse tendency concerning GTS% under Cd treatment. Aly (2012) reported Cd impact in Egyptian clover and Sudan grass seedlings for 21 days. The previous study showed that Cd caused increased P% and decreased GTS% as Cd applied concentration increased. Moreover, Zhang *et al.* (2015) stated Cd effect in two rice varieties by SRAP marker. The previous investigation revealed that GTS% decreased as Cd applied concentration increased. Similarly, Salarizadeh and Reza-Kavousi (2015) reported decreased GTS% under Pb stress using RAPD marker in cumin (*C. cyminum*). Moreover, similar findings were reported in artichoke (*C. scolymus* L.) and runner bean (*P. coccineus* L.) (Candan and Batir, 2015) exposed to Pb stress using same marker. Whereas, Cenkci *et al.* (2010) stated that Pb exposure in fodder turnip (*Brassica rapa* L.) using 11 RAPD primers; caused decline in GTS% as Pb concentration increased from 0.5 to 5 mg/l Pb. Aydin *et al.* (2015) however, reported that GTS% values inversely increased from 78.1 to 90.1% as Pb concentration increased from 40 to 240 mg/l Pb in tomato (*L. esculentum*) after 21 days Pb exposure using RAPD marker.

## CONCLUSION

According to the given reports presented in Table 1, the most investigated pollutants at physiological approach were in the

following order: Cd (71%) > Zn (26%) > Cu (24%) > Pb (19%). From Table 2, it worth noting that the given investigations mainly focused on genotoxicity induced by Cd (57%) followed by Pb (24%), Zn (21%) and Cu (14%). Otherwise, RAPD tool among applied techniques proved to be the most employed technique (80%) as molecular marker for detecting pollutants genotoxicity. Thereby, RAPD marker could be employed as a potent marker for detecting genotoxicity in living organisms.

## ACKNOWLEDGEMENTS

I thank Professor Othman Ibrahim (Director General of AECS) and Professor MirAli Nizar (Head of Molecular Biology and Biotechnology Department )for their support and to the Plant Biotechnology Group for technical assistance.

## REFERENCES

- Aina, R., Labra, M., Fumagalli, P., Vannini, C., Marsoni, M., Cucchi, U., Bracale, M., Sgorbati, S. and Citterio, S. (2007).** Thiol-peptide level and proteomic changes in response to cadmium toxicity in *Oryza sativa* L. roots. *Environmental and Experimental Botany*, **59 (3)**, 381-892.
- Ahmad, M.A., Gaur, R. and Gupta, M. (2012).** Comparative biochemical and RAPD analysis in two varieties of rice (*Oryza sativa*) under arsenic stress by using various biomarkers. *Journal of Hazardous Materials*, **217-218**, 141-148.
- Aksoy,Ö., Deveci, A. and Tütünoğlu, B. (2015).** Genotoxic effects of the fungicide

sportak on zea mays.Fresenius Environmental Bulletin, **24 (2)**, 552-559.

**Al-Khateeb, W. and Al-Qwasemeh, H. (2014)**. Cadmium, copper and zinc toxicity effects on growth, proline content and genetic stability of *Solanum nigrum* L., a crop wild relative for tomato; comparative study. Physiology and Molecular Biology of Plants, **20 (1)**, 31-39.

**Al-Qurainy, F. (2009)**. Toxicity of heavy metals and their molecular detection on *Phaseolus vulgaris* (L.). Australian Journal of Basic and Applied Sciences, **3 (3)**,3025-3035.

**Al-Qurainy, F. (2010)**. Application of inter simple sequence repeat (ISSR marker) to detect genotoxic effect of heavy metals on *Eruca sativa* (L.). African Journal of Biotechnology, **9 (4)**, 467-474.

**Al-Qurainy, F., Alameri, A.A. and Khan, S. (2010)**. RAPD profile for the assessment of genotoxicity on a medicinal plant; *Eruca sativa*. Journal of Medicinal Plant Research, **4 (7)**, 579-586.

**Aly, A.A. (2012)**.Application of DNA (RAPD) and ultrastructure to detect the effect of cadmium stress in Egyptian clover and Sudan grass plantlets.The Journal of Stress Physiology and Biochemistry, **8 (1)**, 241-257.

**Amado Filho, G.M., Karez, C.S., Pfeiffer, W.C., Yoneishigue-Valentin, Y. and Farina, M. (1996)**. Accumulation effects on growth and localisation of zinc in *Padina*

*gymnospora* (Dictyotales, Phaeophyceae). Hydrobiologia, **326/327**, 451–456.

**Amado Filho, G.M., Karez, C.S., Andrade, L.R., Yoneshigue-Valentin, Y. and Pfeiffer, W.C. (1997)**. Effects on growth and accumulation of zinc in six seaweed species. Ecotoxicology and Environmental Safety, **37 (3)**, 223–228.

**Aslam, R., Ansari, M.Y.K., Choudhary, S., Bhat, T.M. and Jahan, N. (2014)**. Genotoxic effects of heavy metal cadmium on growth, biochemical, cyto-physiological parameters and detection of DNA polymorphism by RAPD in *Capsicum annum* L. – An important spice crop of India. Saudi Journal of Biological Sciences, **21**, 465–472.

**Atienzar, F., Evenden, A., Jha, A., Savva, D. and Depledge, M. (2000)**. Optimized RAPD analysis generates high-quality genomic DNA profiles at high annealing temperature. Biotechniques, **28 (1)**, 52-54.

**Atienzar, A., Venier, P., Jha, A.N. and Depledge, M.H. (2002)**. Evaluation of the random amplified polymorphic DNA (RAPD) assay for the detection of DNA damage and mutations. Mutation Research, **521 (1-2)**, 151–163.

**Atienzar, F.A. and Jha, A.N. (2006)**. The random amplified polymorphic DNA (RAPD) assay and related techniques applied to genotoxicity and carcinogenesis studies: a critical review. Mutation Research, **613 (2-3)**, 76-102.

**Aydın, S.S., Gökçe, E., Büyük, İ. and Aras, S. (2012).** Characterization of Copper and Zinc induced stress on Cucumber (*Cucumis sativus* L.) by molecular and population parameters. *Mutation Research*, **746 (1)**, 49-55.

**Aydın, S.S., Başaran, E., Cansaran-Duman, D. and Aras, S.(2013).** Genotoxic effect of cadmium in okra (*Abelmoschus esculantus* L.) seedlings: comparative investigation with population parameter and molecular marker. *Journal of Environmental Biology*,**34**, 985-990.

**Aydın, S.S., Kiliç, Z., Dönmez, Ç., Altunkaynak, E. and Aras, S. (2015).** Assesment of genotoxicity induced by lead pollution in tomato (*Lycopersicum esculentum*) by molecular and population markers. *Biological Diversity and Conservation*, **8 (1)**, 83-89.

**Azimi, A., Shahriari, F., Fotovat, A., Qale, R.K. and Agje, K.H. (2013).** Investigation of DNA changes in wheat (*Triticum aestivum* L.) induced by cadmium using random amplified polymorphic DNA (RAPD) analysis. *African Journal of Biotechnology*, **12(16)**, 1921-1929.

**Bal, W. and Kasprzak, K.S. (2002).** Induction of oxidative DNA damage by carcinogenic metals. *Toxicology letters*,**127 (1-3)**, 55-62.

**Bing, H., Wu, Y., Liu, E. and Yang, X. (2013).** Assessment of heavy metal enrichment and its human impact in lacustrine sediments from four lakes in the mid-low

reaches of the Yangtze River, China. *Journal of Environmental Sciences*, **25 (7)**, 1300-1309.

**Bulgariu, D. and Bulgariu, L. (2012).** Equilibrium and kinetics studies of heavy metal ions biosorption on green algae waste biomass. *Bioresource Technology*, **103 (1)**,489-493.

**Bouzon, Z.L., Ferreira, E.C., dos Santos, R., Scherner, F., Horta, P.A. Maraschin, M. and Schmidt, E.C. (2012).** Influences of cadmium on fine structure and metabolism of *Hypnea musciformis* (Rhodophyta, Gigartinales) cultivated *in vitro*. *Protoplasma*, **249 (3)**, 637–650.

**Caldelas, C., Araus, J.L., Febrero, A. and Bort, J. (2012).** Accumulation and toxic effects of chromium and zinc in *Iris pseudacorus* L. *Acta Physiologiae Plantarum*, **34 (3)**, 1217-1228.

**Cenkci, S., Yildiz, M., Ciğerci, I.H., Konuk, M. and Bozdağ, A. (2009).** Toxic chemicals-induced genotoxicity detected by random amplified polymorphic DNA (RAPD) in bean (*Phaseolus vulgaris* L.) seedlings. *Chemosphere*, **76 (7)**, 900–906.

**Collén, J., Pinto, E., Pedersen, M. and Colepicolo, P. (2003).** Induction of oxidative stress in the red macroalga *Gracilaria tenuistipitata* by pollutants metals. *Archives of Environmental Contamination and Toxicology*, **45**, 337–342.

**dos Santos, R.W., Schmidt, É.C., Martins, R.P., Latini, A., Maraschin, M., Horta, P.A. and Bouzon, Z.L. (2012).** Effects of cadmium on growth, photosynthetic pigments, photosynthetic performance, biochemical parameters and structure of chloroplasts in the Agarophyte *Gracilaria domingensis* (Rhodophyta, Gracilariales). *American Journal of Plant Sciences*, **3 (8)**, 1077-1084.

**Enan, M.R., (2006).** Application of random amplified polymorphic DNA (RAPD) to detect the genotoxic effect of heavy metals. *Biotechnology and Applied Biochemistry*, **43(3)**, 147-154.

**Erturk, F.A., Nardemir, G., Ay, H., Arslan, E. and Agar, G. (2015).** Determination of genotoxic effects of boron and zinc on *Zea mays* using protein and random amplification of polymorphic DNA analyses. *Toxicology and Industrial Health*, **31**, 1015-1023.

**Fusconi, A., Gallo, C. and Wanda, C. (2007).** Effects of cadmium on root apical meristems of *Pisum sativum* L. Cell viability, cell proliferation and microtubule pattern as suitable markers for assessment of stress pollution. *Mutation Research*, **632 (1-2)**, 9-19.

**Ghiani, A., Fumagalli, P., Nguyen-Van, T., Gentili, R. and Citterio, S. (2014).** The combined toxic and genotoxic effects of Cd and As to plant bioindicator *Trifolium repens* L. *PLoS ONE*, **9 (6)**, 1-11.

**Ghnaya, T., Slama, I., Messedi, D., Grignon, C., Ghorbel, M.H. and Abdel, C. (2007).** Effects of Cd<sup>2+</sup> on K<sup>+</sup>, Ca<sup>2+</sup> and N uptake in two halophytes *Sesuvium portulacastrum* and *Mesembryanthemum crystallinum*: Consequences on growth. *Chemosphere*, **67 (1)**, 72-79.

**Gichner, T., Patková, Z., Száková, J. and Demnerová, K. (2004).** Cadmium induces DNA damage in tobacco roots, but no DNA damage, somatic mutations or homologous recombination in tobacco leaves. *Mutation Research*, **559 (1-2)**, 49-57.

**Gupta, M. and Sarin, N.B. (2009).** Heavy metal induced DNA changes in aquatic macrophytes: Random amplified polymorphic DNA analysis and identification of sequence characterized amplified region marker. *Journal of Environmental Science*, **21 (5)**, 686-690.

**Hori, K., Maruyama, F., Fujisawa, T., Togashi, T., Yamamoto, N., Seo, M. et al. (2014).** *Klebsormidium flaccidum* genome reveals primary factors for plant terrestrial adaptation. *Nature Communication*, **5**, 1-9.

**Jamers, A., Blust, R., De-Coena, W., Griffinab, J.L. and Jonesco, O.A.H. (2013).** An omics based assessment of cadmium toxicity in the green alga *Chlamydomonas reinhardtii*. *Aquatic Toxicology*, **126 (1)**, 355-364.

**Jana, S. and Choudhuri, M.A. (1984).** Synergistic effects of heavy metal

pollutants on senescence in submerged aquatic plants. *Water, Air, and Soil Pollution*, **21 (1)**, 351–357.

**Jiang, H., Gao, B., Li, W., Zhu, M., Zheng, C., Zheng, Q. and Wang, C. (2013).** Physiological and biochemical responses of *Ulva prolifera* and *Ulva linza* to cadmium stress. *The Scientific World Journal*, **2013**, 1-11.

**Kaplan, D., Heimer, E Y.M., Abeliovich, A. and Goldsbroug, B. (1995).** Cadmium toxicity and resistance in *Chlorella sp.* *Plant Science*, **109 (2)**, 129-137.

**Kekec, G., Sakcali, M.S. and Uzonur, I. (2010).** Assessment of genotoxic effects of boron on wheat (*Triticum aestivum* L.) and bean (*Phaseolus vulgaris* L.) by using RAPD analysis. *Bulletin of Environmental Contamination and Toxicology*, **84 (6)**, 759–764.

**Khan, M.U., Malik, R.N. and Muhammad, S. (2013).** Human health risk from Heavy metal via food crops consumption with wastewater irrigation practices in Pakistan. *Chemosphere*, **93 (10)**, 2230-2238.

**Körpe, D.A. and Aras, S. (2011).** Evaluation of copper-induced stress on eggplant (*Solanum melongena* L.) seedlings at the molecular and population levels by use of various biomarkers. *Mutation Research*, **719 (1-2)**, 29-34.

**Kupper, H., Kupper, F. and Spiller, M. (1996).** Environmental relevance of heavy metal substituted chlorophylls using the

example of water plants. *Journal of Experimental Botany*, **47 (295)**, 259–266.

**Labra, M., Di Fabio, T., Grassi, F., Regondi, S.M., Bracale, M., Vannini, C. and Agradi, E.(2003).** AFLP analysis as biomarker of exposure to organic and inorganic genotoxic substances in plants. *Chemosphere*, **52 (7)**, 1183-1188.

**Liu, W., Li, P., Qi, X.M., Zhou, Q., Zheng, L., Sun, T. and Yang, Y.(2005).** DNA changes in barley (*Hordeum vulgare*) seedlings induced by cadmium pollution using RAPD analysis. *Chemosphere*, **61 (2)**, 158-167.

**Liu, W., Yang, Y., Zhou, Q., Xie, L., Li, P. and Sun, T.(2007).** Impact assessment of cadmium contamination on rice (*Oryza sativa* L.) seedlings at molecular and population levels using multiple biomarkers. *Chemosphere*, **67 (6)**, 1155-1163.

**Liu, W., Sun, L., Zhong, M., Zhou, Q., Gong, Z., Li, P., Tai, P. and Li, X.(2012).** Cadmium-induced DNA damage and mutations in *Arabidopsis* plantlet shoots identified by DNA fingerprinting. *Chemosphere*, **89 (9)**, 1048–1055.

**Milone, M.T., Sgherri, C., Clijsters, H. and Navari-Izzo, F. (2003).** Antioxidative responses of wheat treated with realistic concentration of cadmium. *Environmental Experimental Botany*, **50 (3)**, 265–276.

**Ortega-Villasante, C., Rellán-Alvarez, R., DelCampo, F.F., Carpena-Ruiz, R.O. and Hernández, L.E. (2005).** Cellular damage induced by cadmium and mercury in

*Medicago sativa*. Journal of Experimental Botany, **56 (418)**, 2239-2251.

**Ova, D. and Övez, B. (2013).** Characterization of heavy metal biosorption from aqueous solutions with *Nitzschia closterium* biomass, Digital Proceeding of the ICOEST'2013 - Cappadocia. C.Ozdemir, S. Şahinkaya, E. Kalıpcı, M.K. Oden (editors), Nevsehir-Turkey, June, pp. 18 –21.

**Ozyigit I.I., Yilmaz S., Dogan I., Sakcali M.S., Tombuloglu G. and Demir G. (2016a).** Detection of physiological and genotoxic damages reflecting toxicity in *Kalanchoe* clones. Global NEST Journal, **18 (1)**, 223-232.

**Ozyigit, I.I., Dogan, I., Igdelioglu, S., Filiz, E., Karadeniz, S. and Uzunova, Z. (2016b).** Screening of damage induced by lead (Pb) in rye (*Secale cereale* L.) – a genetic and physiological approach. Biotechnology and Biotechnological Equipment, **30 (3)**, 489-496.

**Pazoki, A., Ilkaee, M.N., Far, M.D. and Golzardi, F. (2014).** Impact of bio-fertilizers and different levels of lead on pigment content of wheat (*Triticum aestivum* L.). Journal of Biodiversity and Environmental Sciences, **5**, 348-355.

**Phillips, D.J.H. and Rainbow, P.S. (1993).** Biomonitoring of Trace Aquatic Contaminants. CRC Press, Boca Raton, FL.

**Prabha, Y., Soni, S.K., Gupta, S. and Sonal. 2016.** Potential of algae in bioremediation of wastewater: Current research. International Journal of Current Microbiology and Applied Sciences, **5 (2)**, 693-700.

**Rainbow, P.S. (1995).** Biomonitoring of heavy metal availability in the marine environment. Marine Pollution Bulletin, **31**, 183–192.

**Raj, A., Kumar, S., Haq, I. and Kumar, M. (2014).** Detection of tannery effluents induced DNA damage in mung bean by use of Random Amplified Polymorphic DNA markers. ISRN Biotechnology, **2014**, 1-8.

**Salarizadeh, S. and Reza-Kavousi, H. (2015).** Application of random amplified polymorphic DNA (RAPD) to detect the genotoxic effect of cadmium on two Iranian ecotypes of cumin (*Cuminum cyminum*). Journal of Cell and Molecular Research, **7 (1)**, 38-46.

**Saleh, B. (2015a).** Physiological response of the green algae *Ulva lactuca* (Chlorophyta) to heavy metals stress. Journal of Stress Physiology & Biochemistry, **11 (3)**, 38-51.

**Saleh, B. (2015b).** Detection of genetic variations in marine algae *Ulva lactuca* (Chlorophyta) induced by heavy metal pollutants. The Journal of Stress Physiology and Biochemistry, **11 (3)**, 26-37.

**Saleh, B. (2016a).** Genomic DNA changes in *Ulva lactuca* (Chlorophyta) under heavy



metal stress. International Journal of Environmental Sciences **7 (3)**, 245-255.

**Saleh, B. (2016b).** Cadmium stress assessment in *Ulva lactuca* (Chlorophyta) and *Padina pavonica* (Phaeophyta) marine algae. Journal of Bioinnovation, **5 (6)**, 835-849.

**Saleh, B. (2016c).** Genetic variation in *Ulva lactuca* (Chlorophyceae) marine algae under cadmium stress. Journal of Bioinnovation, **5 (5)**, 673-685.

**Saleh, B. (2016d).** DNA alteration induced by cadmium (Cd) metal in *Padina pavonica* (Phaeophyta) marine brown algae. Plant Cell Biotechnology and Molecular Biology, **17 (7-8)** (In press).

**Saleh, B. (2016e).** Lead (Pb) heavy metal impacts in the green *Ulva lactuca* (Chlorophyceae) marine algae. The Journal of Stress Physiology & Biochemistry, **12 (2)**, 62-71.

**Saleh, B. and Al-Mariri, A. 2016.** Antimicrobial activity of the marine algal extracts against selected pathogens. Journal of Agricultural Science and Technology (In Press).

**Shah, K., Ritambhara, G.K., Verma, S. and Dubey, R.S. (2001).** Effect of cadmium on lipid peroxidation, superoxide anion generation and activities of antioxidant enzymes in growing rice seedlings. Plant Science. **161 (6)**, 1135–1144.

**Shahrtash, M., Mohsenzadeh, S. and Mohabatkar, H. (2010).** Cadmium-induced

genotoxicity detected by the random amplification of polymorphism DNA in the maize seedling roots. Journal of Cell and Molecular Research, **2 (1)**, 42-48.

**Shariati, M. and Yahyaabadi, S. (2006).** The effect of different concentrations of cadmium on the growth rate and beta-carotene synthesis in unicellular green algae *Dunaliella salina*. Iranian Journal of Science and Technology, **30(A1)**, 57-63.

**Trevors, J.T., Stratton, G.W. and Gadd, G.M. (1986).** Cadmium transport, resistance, and toxicity in bacteria, algae, and fungi. Canadian Journal of Microbiology, **32 (6)**, 447-464.

**Unal, D., Isik, N.O. and Sukatar, A. (2010).** Effects of chromium VI stress on green alga *Ulva lactuca* (L.). Turkish Journal of Biology, **34**, 119-124.

**Valavanidis, A. and Vlachogianni, T. (2010).** Metal pollution in Ecosystems. Ecotoxicology studies and risk assessment in the marine environment. Science advances on Environment, Toxicology and Ecotoxicology, 1-14.

**Verma, S. and Dubey, R. (2001).** Effect of cadmium on soluble sugars and enzymes of their metabolism in rice. Biologia Plantarum, **44 (1)**, 117-123.

**Vilar, V.J.P., Botelho, C.M.S. and Boaventura, R.A.R. (2008).** Effect of Cu(II), Cd(II) and Zn(II) on Pb(II) biosorption by algae *Gelidium*-derived materials. Journal of Hazardous Materials **154**, 711–720.

**Wang, J.S., Chou, H.N., Fan, J.J. and Chen, C.M. (1998).** Uptake and transfer of high PCB concentrations from phytoplankton to aquatic biota. *Chemosphere*, **36 (6)**, 1201–1210.

**Zhang, X., Chen, H., Jiang, H., Lu, W., Pan, J., Qian, Q. and Xue, D. (2015).** Measuring the damage of heavy metal cadmium in rice seedlings by SRAP analysis combined with physiological and biochemical parameters. *The Journal of the Science of Food and Agriculture*, **95 (11)**, 2292-2298.

**Zengin, F. (2013).** Physiological behavior of bean (*Phaseolus vulgaris* L.) seedlings under metal stress. *Biological Research*, **46(1)**, 79-85.

**Zhou, L., Li, J., Lin, X. and Al-Rasheid, K.A. (2011).** Use of RAPD to detect DNA damage induced by nitrofurazone in marine ciliate, *Euplotes vannus* (Protozoa, Ciliophora). *Aquatic Toxicology*, **103 (3-4)**, 225-232.