

<https://doi.org/10.46344/JBINO.2023.v12i02.14>

## SAFE TREATMENT OF URINARY TRACT INFECTIONS BY AMERICAN CRANBERRY

Muhammad Arshad Ullah<sup>1</sup> and Ali Hassan<sup>2</sup>

<sup>1</sup>Pakistan Agricultural Research Council, Islamabad, Pakistan

<sup>2</sup>PMAS- University of Arid Agriculture, Rawalpindi, Pakistan

### ABSTRACT

Cranberry fruit (*Vaccinium macrocarpon*) grows on evergreen shrubs that are native to North America. Cranberry is a term derived from the contraction of "crane berry." This name is derived from the nickname of the bilberry flower, which, when it withers, is similar in appearance to the head and neck of the sand crane, a bird that often feeds on the berries of this plant. The cranberry is part of the Ericaceae family and naturally grows in acidic swamps full of peat moss in humid forests. Cranberries are composed of water (88%), organic acids (including salicylate), fructose, vitamin C (high levels, i.e., 200 mg/kg of fresh berries), flavonoids, anthocyanidins, catechins, and triterpenoids. The chemical constituents responsible for their taste are the iridoid glycosides. The anthocyanidins and proanthocyanidins (PAC) are tannins (stable polyphenols) found only in vaccinium berries and function as a natural plant defense system against microbes. Common preparations with cranberries include fresh, whole berries, gelatinized products, juices (usually 10-25% pure juice) and capsules. Pure juice is too acidic (pH, 2.5) and unpalatable, even with sweeteners. Despite cranberry presentation, it is generally recommended to consume cranberries just prior or two hours after meals; it is also important to drink lots of water, mainly after preparations from dehydrated juices. Cranberry juice, predominantly in the form of a juice cocktail drink with approximately 25% cranberry juice, has been the traditional choice of most women seeking to prevent Urinary Tract Infections (UTIs). American cranberry has a complex and rich phytochemical composition, particularly flavan-3-ols, A-type procyanidins (PACs), anthocyanins, benzoic acid, and ursolic acid. Cranberry flavan-3-ols are present as monomers, oligomers, and polymers). These oligomers and polymers are also referred to as PACs or condensed tannins and represent 85% of the total flavan-3-ols on a weight basis. Cranberry is the main source of peonidin among 100 foods commonly consumed in the United States. Quercetin 3-galactoside is the predominant form, but at least 11 other glycosides are present in lower concentrations. Some of these, such as quercetin-3-acetylramnoside are rare in berries. In the Phenol Explorer database, the flavonol content of plant foods is usually <3 mg/100 g FW, although bilberry, black-berry, and blueberry contain 3.2–17 mg/kg. Cranberry fruit is classed as a functional food due to the naturally high content of compounds, such as polyphenols, which are believed to have antioxidant and therefore health-promoting properties. Health benefits of cranberry consumption range from cardio protective effects due to improved cholesterol profiles to aiding digestive health. Cranberry exists in various forms, including the raw fruit (fresh and dried), cranberry juice and

cranberry extract in capsule/tablet formulations. Cranberry extract could be a potential alternative to antibiotics to treat acute uncomplicated UTIs. Proanthocyanidin (PAC) with A-type linkages, or their metabolites, is believed to be the active ingredient in cranberry, preventing *Escherichia coli* (*E. coli*) from binding to the bladder uroepithelium and thereby reducing the ability of *E. coli* to cause and sustain a UTI. Cranberries have also been found to improve lipid profile, improve endothelial function, and lower several markers of cardio metabolic risk. Nowadays, growing evidence suggests an important role of cranberries in maintaining digestive health. In addition to the anti-inflammatory effects, cranberries may also influence intestinal barrier integrity, which is another essential element of intestinal health. Cranberry was reported as the main source of peonidin among 100 foods commonly consumed in the United States. However, in the majority of studies, the total anthocyanin content is re-ported rather than amounts of individual anthocyanins. This approach may change because the bioavailability and health effects of anthocyanins seem to be affected by the structures of the aglycones or the glycosidic moieties. Quercetin 3-galactoside is the predominant form, but at least 11 other glycosides are present in lower concentrations. Some of these, such as quercetin-3-acetylramnoside are rare in berries. As shown in the Phenol Explorer database, the flavonol content of plant foods is usually <3 mg/100 g FW, although bilberry, black-berry, and blueberry contain 3.2–17 mg/kg.

## INTRODUCTION

Cranberry is a term derived from the contraction of "crane berry." This name is derived from the nickname of the bilberry flower, which, when it withers, is similar in appearance to the head and neck of the sand crane, a bird that often feeds on the berries of this plant (Guay, 2009). The cranberry is part of the Ericaceae family and naturally grows in acidic swamps full of peat moss in humid forests (Bruyere, 2006). The American cranberry (*Vaccinium macrocarpon*) was historically used by North American Indians to treat UTIs (Guay, 2009). There are other relatives of the cranberry family (European cranberry – *V. oxycoccus*; lingonberry – *V. vitisidaea*; blueberry – *V. myrtillus*) that share some of the cranberry's basic components, but research evidence for a role in prevention is limited (Kontiokari *et al.*, 2001 and Jepson and Craig, 2007).

Cranberries are composed of water (88%), organic acids (including salicylate), fructose, vitamin C (high levels, i.e., 200 mg/kg of fresh berries), flavonoids, anthocyanidins, catechins, and triterpenoids (Guay, 2009). The chemical constituents responsible for their taste are the iridoid glycosides. The anthocyanidins and proanthocyanidins (PAC) are tannins (stable polyphenols) found only in *Vaccinium* berries and function as a natural plant defense system against

microbes (Guay, 2009 and Cimolai and Cimolai, 2007).

Common preparations with cranberries include fresh, whole berries, gelatinized products, juices (usually 10-25% pure juice) and capsules (Guay, 2009; Bruyere, 2006; Kontiokari *et al.*, 2001; Jepson and Craig, 2007; Gupta *et al.*, 2007 and Di Martino *et al.*, 2006). Pure juice is too acidic (pH, 2.5) and unpalatable, even with sweeteners (Guay, 2009). Despite cranberry presentation, it is generally recommended to consume cranberries just prior or two hours after meals; it is also important to drink lots of water, mainly after preparations from dehydrated juices (Bruyere, 2006). Cranberry juice, predominantly in the form of a juice cocktail drink with approximately 25% cranberry juice, has been the traditional choice of most women seeking to prevent Urinary Tract Infections (UTIs). Cranberries have been tested for their clinical relevance in many different conditions. They have been evaluated in the treatment of UTIs but were deemed ineffective (Cimolai and Cimolai, 2007 and Lavigne *et al.*, 2007).

Cranberries were also studied for UTI prophylaxis chiefly in women, but also in children and men; additionally, they have been studied in conditions such as neurogenic bladder and pregnancy, (Kontiokari *et al.*, 2001; Stothers, 2002; Barbosa-Cesnik *et al.*, 2011; McMurdo *et al.*, 2005; 2009; Wing *et al.*, 2008; Lee *et al.*, 2007; McGuinness *et al.*, 2002; Waites *et al.*, 2004 and Ferrara *et al.*, 2008). Most clinical interest in the use of cranberries is for

cystitis prevention. In the 2008 Cochrane Database of Systematic Reviews, there were ten randomized trials regarding UTI prevention on a total of 1,049 patients. They concluded that there is some evidence that cranberry juice may decrease the number of symptomatic UTIs over a 12 month period, particularly, and only, for women with recurrent UTIs (Jepson and Craig, 2008). Whereas berries are noted simply as good sources of potassium or fiber, recent research suggests that berry fruits are a rich source of numerous phytochemicals with a broad array of bioactivity and an impact on human health (Paredes-López et al., 2010; Côté et al., 2010; Basu and Lyons, 2012; Wang et al., 2012 and Kaspar and Khoo, 2013).

The American cranberry (*Vaccinium macrocarpon*) is a particularly rich source of (poly)phenols, which have been associated in vitro with antibacterial, antiviral, antimutagenic, anticarcinogenic, antitumorigenic, antiangiogenic, anti-inflammatory, and antioxidant properties (Côté et al., 2010; McKay and Blumberg, 2007 and Del Rio et al., 2013). In vivo, animal models reveal that cranberry extracts can reduce C-reactive protein (CRP) and proinflammatory interleukins and increase NO synthesis (Kim et al., 2011); decrease angiotensin-converting enzyme, angiotensin II, and angiotensin II type 1 receptor (Yung et al., 2011); suppress *Helicobacter pylori* infection (Xiao and Shi, 2003); and improve pancreatic  $\beta$ -cell glucose responsiveness and functional  $\beta$ -cell mass (Zhu et al., 2011). Some of these

actions may underlie the results from clinical studies showing that cranberry products can lower LDL cholesterol (LDL-C) and total cholesterol (Lee et al., 2008), increase HDL cholesterol (HDL-C) while lowering the oxidative modification of LDL-C (Ruel et al., 2006), improve endothelial function (Dohadwala et al., 2011 and Flammer et al., 2011), lower glycemic responses (Wilson et al., 2010), elevate plasma antioxidant capacity (Ruel et al., 2005; Duthie et al., 2006 and Vinson et al., 2008), modulate ulcerogenic gastric *H. pylori* colonization (Gotteland et al., 2008 and Shmueli et al., 2007), decrease cariogenic *Streptococcus mutans* and total bacterial counts in saliva (Weiss et al., 2004), reduce biomarkers of metabolic syndrome (Basu and Lyons, 2012 and Basu et al., 2011), and protect against urinary tract infections (UTIs) (Wang et al., 2012 and Vasileiou et al., 2013).

American cranberry has a complex and rich phytochemical composition, particularly flavan-3-ols, A-type procyanidins (PACs), anthocyanins, benzoic acid, and ursolic acid. Cranberry flavan-3-ols are present as monomers, oligomers, and polymers (Pappas and Schaich, 2009). These oligomers and polymers are also referred to as PACs or condensed tannins and represent 85% of the total flavan-3-ols on a weight basis (White et al., 2011 and Gu et al., 2004).

Cranberry was reported as the main source of peonidin among 100 foods commonly consumed in the United States (Wu et al., 2006). However, in the majority

of studies, the total anthocyanin content is reported rather than amounts of individual anthocyanins. This approach may change because the bioavailability and health effects of anthocyanins seem to be affected by the structures of the aglycones or the glycosidic moieties (Crozier et al., 2010 and Czank et al., 2013). Quercetin-3-galactoside is the predominant form, but at least 11 other glycosides are present in lower concentrations (Pappas and Schaich, 2009 and Côté et al., 2008). Some of these, such as quercetin-3-acetyl rhamnoside are rare in berries (Mikulic-Petkovsek et al., 2012). As shown in the Phenol Explorer database, the flavonol content of plant foods is usually <3 mg/100 g FW, although bilberry, blackberry, and blueberry contain 3.2–17 mg/kg (Mikulic-Petkovsek et al., 2012 and Harnly et al., 2006).

The other species are blueberry (*Vaccinium angustifolia*) and bilberry (*Vaccinium myrtillus*). Cranberry typically grows in bogs and is a member of the same family as blueberry and bilberry. Massachusetts and Wisconsin are the main areas of present-day commercial production of cranberry (Nova Scotia Department of Agriculture and Fisheries, 2004). Cranberry has been found to specifically inhibit hemagglutination of *E. coli* by expression of types (Nova Scotia Department of Agriculture and Fisheries, 2004) and P adhesion through the component compounds fructose and proanthocyanidins. In the United States, one of every five women has been reported to have a lifetime incidence of

UTI. Of these women, 3 percent experience recurrent disease. Eleven million women receive medication for UTIs annually (Foxman et al., 2000). A recent Cochrane Database systematic review Jepson et al. (2004) found no randomized trials assessing the effectiveness of cranberry juice in the treatment of UTIs and concluded that there is no evidence to support its use. There is much greater evidence-based information available for the use of cranberry in UTI prophylaxis. A Cochrane Database systematic review, Jepson et al. (2004) citing small sample sizes and the poor quality of available trials, determined that there was no reliable evidence of effectiveness of cranberry in UTI prophylaxis.

However, since 2001, two good-quality studies have been published. The first trial of 150 women consisted of three arms: (Nova Scotia Department of Agriculture and Fisheries, 2004) cranberry/lingon berry juice, pro-biotic supplementation with *Lactobacillus GG* drink; and no intervention for 12 months. Findings were a statistically significant 20 percent reduction in absolute risk of infection in women receiving cranberry (number needed to treat: 5) compared with no effect in the probiotic-supplementation and no-intervention groups (Kontiokari et al., 2001). Most recently, a randomized, placebo-controlled trial, a single experimental study showed that the "high-molecular-weight constituent" of cranberry juice that inhibits the adherence of *E. coli* was effective in reversing and inhibiting the coaggregation

of a large portion of dental plaque bacteria. Cranberry also has been recommended as an adjunctive treatment for *Candida* infections (Stothers, 2002). A small study found a significant rise in urinary oxalate levels, prompting a caution that regular use of cranberry may increase the risk of kidney stone formation in patients with a history of oxalate calculi (Terris et al., 2001).

Cranberries have historically been associated with urinary tract health, particularly among women with rUTIs (Raz et al., 2004; Guay, 2009 and Pérez-López et al., 2009). Results from several clinical studies have suggested that cranberries may decrease rUTIs in healthy women (Guay, 2009; Vasileiou et al., 2013; Caljouw et al., 2014; Stapleton et al., 2012 and Takahashi et al., 2013). In addition, in vitro and ex vivo research has suggested that cranberry-derived compounds such as A-type proanthocyanidins and other polyphenols may interfere with adhesion of bacteria (including multidrug-resistant *Escherichia coli*) to epithelial cells of the urinary tract, attenuate the development of uropathogen reservoirs (i.e., in the gastrointestinal tract and intracellular pods within the urothelium), and suppress inflammatory cascades (Vasileiou et al., 2013; Gupta et al., 2012 and Blumberg et al., 2013).

Specifically, a meta-analysis by Wang and colleagues published in 2012 concluded that "cranberry products were associated with protective effects against UTIs (RR: 0.62; 95% CI: 0.49, 0.80), particularly for women with Recurring Urinary Tract

Infections (rUTIs) (RR: 0.53; 95% CI: 0.33, 0.83)" (Wang et al., 2012). In contrast, a meta-analysis by the Cochrane Collaboration, also published in 2012, concluded that ".cranberry juice is less effective than previously indicated. Cranberry juice cannot currently be recommended for the prevention of UTIs" (Jepson et al., 2012). It is interesting that the Cochrane analysis was an up-date of a 2008 report that resulted in a conclusion similar to that derived by Wang et al., indicating a shift in the conclusions from this group (Jepson et al., 2008 and 2012). In theory, meta-analysis of results from randomized clinical trials examines the consistency of data across studies and is considered to be the strongest level of evidence that guides relevant practice decisions (Melnyk, 2004).

Cranberry fruit (*Vaccinium macrocarpon*) grows on evergreen shrubs that are native to North America (Polashock et al., 2014). Cranberry fruit is classed as a functional food due to the naturally high content of compounds, such as polyphenols, which are believed to have antioxidant and therefore health-promoting properties (Szajdek and Borowska, 2008). The reported health benefits of cranberry consumption range from cardio protective effects due to improved cholesterol profiles (Ruel et al., 2006) to aiding digestive health (Pappas et al., 2009). Cranberry exists in various forms, including the raw fruit (fresh and dried), cranberry juice and cranberry extract in capsule/tablet formulations (Bodet et al., 2008). Cranberry extract could be a potential alternative to

antibiotics to treat acute uncomplicated UTIs. Proanthocyanidin (PAC) with A-type linkages, or their metabolites, is believed to be the active ingredient in cranberry, preventing *Escherichia coli* (*E. coli*) from binding to the bladder uroepithelium (Howell et al., 2005) and thereby reducing the ability of *E. coli* to cause and sustain a UTI. Systematic reviews assessing the use of cranberry in the management of recurrent UTIs provide mixed evidence for benefit (Wang et al., 2012 and Luís et al., 2017). A 2012 Cochrane review of 24 trials (n = 4473) of men, women and children found that cranberry did not significantly reduce recurrent UTI compared with placebo, advice to increase water intake or no treatment. A subgroup analysis of women with recurrent UTI found that cranberry consumption resulted in a non-significant reduction in recurrent UTIs (Jepson et al., 2012). Whilst many studies have evaluated the effectiveness of cranberry extract in reducing recurrent UTI, few have assessed effects on symptoms of acute UTIs (Vicariotto, 2014). Cranberries for the prevention of UTIs in susceptible populations are examined in another review by the same authors (Jepson and Craig, 2008).

Cranberries are among a few foods that contain A-type proanthocyanidins. Unlike B-type proanthocyanidins, A-type proanthocyanidins have an additional ether interflavan bond between C2→O→C7 (Prior and Gu, 2005). It was suggested that A-type proanthocyanidins have greater bioactivity compared to B-type (Howell et al., 2005). Two recent

studies suggested that cranberry oligosaccharides might also play a role in preventing UTIs because xyloglucan oligosaccharides were found to be a new cranberry bioactive component with *E. coli* anti-adhesion] activity (Hotchkiss et al., 2015 and Sun et al., 2015).

Cranberries have also been found to improve lipid profile, improve endothelial function, and lower several markers of cardio metabolic risk (Paquette et al., 2017; Lee et al., 2008 and Novotny et al., 2015). Nowadays, growing evidence suggests an important role of cranberries in maintaining digestive health (Anhe et al., 2015 and Denis et al., 2015). Two 250 mL bottles of cranberry juice daily for 90 days was found to suppress *H. Pylori* infection in the stomach of susceptible population (Zhang et al., 2005). In addition to the anti-inflammatory effects, cranberries may also influence intestinal barrier integrity, which is another essential element of intestinal health (Contreras et al., 2015 and Pierre et al., 2014).

In the early 2000s, Kontiokari et al. (2001) and Stothers et al. (2002) investigated the effects of cranberry juice or tablets, respectively, on UTI prevention. The subjects were healthy, sexually active women, aged 21–72 years, with a history of UTI. Both studies showed a significantly decreased number of patients experiencing at least one episode of UTI per year. In a very recent randomized, double-blind, placebo-controlled, multicenter clinical trial, 28373 healthy women with a recent UTI history were recruited to consume a 240

mL low-calorie cranberry juice cocktail daily for 24 weeks. Cranberry juice consumption significantly reduced the number of UTIs by 39%.

Urinary tract infection is a common bacterial infection in children (Habib, 2012). One serving of 50 mL cranberry juice was given to the children to drink once per day for 6 months (Ferrara et al., 2009). This trial had a relatively large sample size (n=84) and low dropout rate, 3.5% in the cranberry group versus 6.8% in the control group. Cranberry juice, with or without proanthocyanidins, was randomly assigned to 40 children for one year (Afshar et al., 2012). A 65% reduction in the risk of UTIs was obtained during a 12-month follow up. The compliance level was 70% for both groups. Both studies suggested that cranberry juice was effective in preventing the recurrence of pediatric UTI. Salo et al. (2012) conducted a double-blinded randomized placebo-controlled trial in 2012.

A 2008 Cochrane review, (Jepson and Craig, 2008) which included five cross-over and five parallel designed studies, reported that cranberry products significantly reduced the incidence of UTIs at 12 months compared to the placebo/control group. Cranberry-containing products were found to be associated with preventative effects on recurrent UTIs in another systematic review of 13 randomized controlled trials (Wang et al., 2012). The most recent meta-analysis and trial sequential analysis evaluated 28 clinical trials and concluded that the use of cranberry products significantly reduced

the incidence of UTIs (Luis et al., 2017). However, meta-analysis in the 2012 Cochrane review<sup>36</sup> concluded that cranberry products did not significantly reduce the occurrence of symptomatic UTI overall compared to placebo, water, or no treatment. This review had substantial heterogeneity in the results.

Several factors contribute to the conflicting results obtained from clinical trials. The major factors are low compliance due to the asstringency and bitterness of cranberry juice, high withdrawal rate, inconsistency of dosage, and lack of standardization of cranberry content in testing products. Cranberry capsules were used to overcome the undesired taste of cranberry juice. However, in two trials using cranberry capsules, the withdrawal rates were more than 40%, partly due to the side effects (Linsenmeyer et al., 2004 and Waites et al., 2004). Quite a few trials were not randomized, double blind, or placebo controlled, or had short duration time (less than 6 months). In addition, the optimum dosage and formulation were not established in previous studies (Guay, 2004).

## REFERENCES

- Afshar K, Stothers L, Scott H and MacNeily AE, Cranberry Juice for the prevention of pediatric urinary tract infection: a randomized controlled trial. *JUrol*188:1584–1587 (2012).
- Anhe FF, Roy D, Pilon G, Dudonne S, Matamoros S, Varin TV et al., A polyphenol-rich cranberry extract protects from diet-induced obesity, insulin resistance and intestinal inflammation in association with



- increased Akkermansia spp. population in the gut microbiota of mice. *Gut* 64:872–883 (2015).
- Barbosa-Cesnik C, Brown MB, Buxton M, Zhang L, DeBusscher J, Foxman B. Cranberry juice fails to prevent recurrent urinary tract infection: results from a randomized placebo-controlled trial. *Clin Infect Dis*. 2011; 52(1):23-30, <http://dx.doi.org/10.1093/cid/ciq073>.
- Basu A, Betts NM, Ortiz J, Simmons B, Wu M, Lyons TJ. Low-energy cranberry juice decreases lipid oxidation and increases plasma antioxidant capacity in women with metabolic syndrome. *Nutr Res*. 2011;31:190–6.
- Basu A, Lyons TJ. Strawberries, blueberries, and cranberries in the met-abolic syndrome: clinical perspectives. *J Agric Food Chem*. 2012;60: 5687–92.
- Blumberg JB, Camesano TA, Cassidy A, Krist-Etherton P, Howell A, Manach C, Ostertag LM, Sies H, Skulas-Ray A, Vita JA. Cranberries and their bioactive constituents in human health. *Adv Nutr* 2013;4: 618–32.
- Bodet, C.; Grenier, D.; Chandad, F.; Ofek, I.; Steinberg, D.; Weiss, E. Potential oral health benefits of cranberry. *Crit. Rev. Food Sci. Nutr*. 2008, 48, 672–680. [CrossRef] [PubMed]
- Bruyere F. [Use of cranberry in chronic urinary tract infections]. *Med Mal Infect*. 2006; 36(7):358-63, <http://dx.doi.org/10.1016/j.medmal.2006.05.001>.
- Caljouw MA, van den Hout WB, Putter H, Achterberg WP, Cools HJ, Gussekloo J. Effectiveness of cranberry capsules to prevent urinary tract infections in vulnerable older persons: a double-blind randomized placebo-controlled trial in long-term care facilities. *J Am Geriatr Soc* 2014; 62:103–10.
- Cimolai N, Cimolai T. The cranberry and the urinary tract. *Eur J Clin Microbiol Infect Dis*. 2007; 26(11):767-76, <http://dx.doi.org/10.1007/s10096-007-0379-0>.
- Contreras TC, Ricciardi E, Cremonini E and Oteiza PI, (-)-Epicatechin in the prevention of tumor necrosis alpha-induced loss of Caco-2 cell barrier integrity. *Arch Biochem Biophys* 573:84–91 (2015).
- Côté J, Caillet S, Doyon G, Sylvain JF, Lacroix M. Analyzing cranberry bioactive compounds. *Crit Rev Food Sci Nutr*. 2010;50:872–88.
- Crozier A, Del Rio D, Clifford MN. Bioavailability of dietary flavo-noids and phenolic compounds. *Mol Aspects Med*. 2010;31:446–67.
- Czank C, Cassidy A, Zhang Q, Morrison DJ, Preston T, Kroon PA, Botting NP, Kay CD. Human metabolism and excretion of the antho-cyanin, cyanidin-3-glucoside: a 13C-tracer study. *Am J Clin Nutr*. 2013;97:995–1003.
- Del Rio D, Rodriguez-Mateos A, Spencer JP, Tognolini M, Borges G, Crozier A. Dietary (poly)phenolics in human health: structures, bio-availability, and evidence of protective effects against chronic diseases. *Antioxid Redox Signal*. 2013;18:1818–92.
- Denis MC, Desjardins Y, Furtos A, Marcil V, Dudonne S, Montoudis A et al., Prevention of oxidative stress, inflammation and mitochondria-dial dysfunction in the intestine by different cranberry phenolic

- fractions. Clin Sci (Lond) 128:197–212 (2015).
- Di Martino P, Agniel R, David K, Templer C, Gaillard JL, Denys P, et al. Reduction of Escherichia coli adherence to uroepithelial bladder cells after consumption of cranberry juice: a double-blind randomized placebo-controlled cross-over trial. World J Urol. 2006; 24(1):21-7, <http://dx.doi.org/10.1007/s00345-005-0045-z>.
- Dohadwala MM, Holbrook M, Hamburg NM, Shenouda SM, Chung WB, Titas M, Kluge MA, Wang N, Palmisano J, Milbury PE, et al. Effects of cranberry juice consumption on vascular function in patients with coronary artery disease. Am J Clin Nutr. 2011;93:934–40.
- Duthie SJ, Jenkinson AM, Crozier A, Mullen W, Pirie L, Kyle J, Yap LS, Christen P, Duthie GG. The effects of cranberry juice consumption on antioxidant status and biomarkers relating to heart disease and cancer in healthy human volunteers. Eur J Nutr. 2006;45:113–22.
- Ferrara P, Romaniello L, Vitelli O, Gatto A, Serva M, Cataldi L. Cranberry juice for the prevention of recurrent urinary tract infections: a randomized controlled trial in children. Scand J Urol Nephrol. 2009; 43(5):369-72, <http://dx.doi.org/10.3109/00365590902936698>.
- Flammer AJ, Martin EA, Gossel M, Widmer RJ, Lennon RJ, Sexton JA, Loeffler D, Khosla S, Lerman LO, Lerman A. Polyphenol-rich cran-berry juice has a neutral effect on endothelial function but decreases the fraction of osteocalcin-expressing endothelial progenitor cells. Eur J Nutr. 2013;52:289–96.
- Foxman B, Barlow R, D'Arcy H, Gillespie B, Sobel JD. Urinary tract infection: self-reported incidence and associated costs. Ann Epidemiol 2000; 10:509-15.
- Gotteland M, Andrews M, Toledo M, Muñoz L, Caceres P, Anziani A, Wittig E, Speisky H, Salazar G. Modulation of Helicobacter pylori colonization with cranberry juice and Lactobacillus johnsoniiLa1 in children. Nutrition. 2008;24:421–6.
- Gu L, Kelm MA, Hammerstone JF, Beecher G, Holden J, Haytowitz D, Gebhardt S, Prior RL. Concentrations of proanthocyanidins in common foods and estimations of normal consumption. J Nutr. 2004; 134:613–7.
- Guay DR. Cranberry and urinary tract infections. Drugs. 2009; 69(7):775-807, <http://dx.doi.org/10.2165/00003495-200969070-00002>.
- Gupta A, Dwivedi M, Mahdi AA, Nagana Gowda GA, Khetrapal CL, Bhandari M. Inhibition of adherence of multi-drug resistant E. Coli by proanthocyanidin. Urol Res 2012;40:143–50.
- Gupta K, Chou MY, Howell A, Wobbe C, Grady R, Stapleton AE. Cranberry products inhibit adherence of p-fimbriated Escherichia coli to primary cultured bladder and vaginal epithelial cells. J Urol. 2007; 177(6):2357-60, <http://dx.doi.org/10.1016/j.juro.2007.01.114>.
- Habib S, Highlights for management of a child with a urinary tract infection. Int J Pediatr 2012;943653 (2012).
- Harnly JM, Doherty RF, Beecher GR, Holden JM, Haytowitz DB, Bhagwat S, Gebhardt S. Flavonoid content of US fruits, vegetables, and nuts. J Agric Food Chem. 2006; 54:9966–77.

- Hotchkiss AT, Jr., Nunez A, Strahan GD, Chau HK, White AK, Marais JP et al., Cranberry xyloglucan structure and inhibition of Escherichia coli adhesion to epithelial cells. *J Agric Food Chem* 63:5622–5633(2015).
- Howell AB, Reed JD, Krueger CG, Winterbottom R, Cunningham DG and Leahy M, A-type cranberry proanthocyanidins and uropathogenic bacterial anti-adhesion activity. *Phytochemistry* 66:2281–2291 (2005).
- Jepson RG, Craig JC. A systematic review of the evidence for cranberries and blueberries in UTI prevention. *Mol Nutr Food Res.* 2007; 51(6):738-45, <http://dx.doi.org/10.1002/mnfr.200600275>.
- Jepson RG, Mihaljevic L, Craig J. Cranberries for treating urinary tract infections. *Cochrane Database Syst Rev* 2004 ;(3):CD001322.
- Jepson RG, Williams G, Craig JG. Cranberries for preventing urinary tract infections [review]. *Cochrane Database Syst Rev* 2012; 10:CD001321.
- Jepson RG, Craig JC. Cranberries for preventing urinary tract infections. *Cochrane Database of Systematic Reviews* 2008, Issue 1. Art. No: CD001321. [DOI: 10.1002/14651858.CD001321.pub3]
- Kaspar KL, Khoo C. Cranberry polyphenols in the promotion of urinary tract, cardiovascular and emerging health areas. In: Skinner M, Hunter D, editors. *Bioactives in fruit: health benefits and functional foods*. New York: Wiley Blackwell; 2013. p. 273–92.
- Kim MJ, Ohn J, Kim JH, Kwak HK. Effects of freeze-dried cranberry powder on serum lipids and inflammatory markers in lipopolysaccharide treated rats fed an atherogenic diet. *Nutr Res Pract.* 2011;5: 404–11.
- Kontiokari T, Sundqvist K, Nuutinen M, Pokka T, Koskela M, Uhari M. Randomised trial of cranberry-lingonberry juice and Lactobacillus GG drink for the prevention of urinary tract infections in women. *BMJ.* 2001; 322(7302):1571, <http://dx.doi.org/10.1136/bmj.322.7302.1571>.
- Lavigne JP, Bourg G, Combescure C, Botto H, Sotto A. In-vitro and in-vivo evidence of dose-dependent decrease of uropathogenic Escherichia coli virulence after consumption of commercial Vaccinium macrocarpon (cranberry) capsules. *Clin Microbiol Infect.* 2008;14(4):350-5, <http://dx.doi.org/10.1111/j.1469-0691.2007.01917.x>.
- Lee BB, Haran MJ, Hunt LM, Simpson JM, Marial O, Rutkowski SB, et al. Spinal-injured neuropathic bladder antiseptics (SINBA) trial. *Spinal Cord.* 2007; 45(8):542-50, <http://dx.doi.org/10.1038/sj.sc.3101974>.
- Lee IT, Chan YC, Lin CW, Lee WJ and Sheu WHH, Effect of cranberry extracts on lipid profiles in subjects with Type 2 diabetes. *Diabetic Med* 25:1473–1477 (2008).
- Linsenmeyer TA, Harrison B, Oakley A, Kirshblum S, Stock JA and Millis SR, Evaluation of cranberry supplement for reduction of urinary tract infections in individuals with neurogenic bladders secondary to spinal cord injury. A prospective, double-blinded, placebo-

- controlled, cross over study. *J Spinal Cord Med* 27:29–34 (2004).
- Luís, Â.; Domingues, F.; Pereira, L. Can cranberries contribute to reduce the incidence of urinary tract infections? A systematic review with meta-analysis and trial sequential analysis of clinical trials. *J. Urol.* 2017, 198, 614–621.
- Maki KC, Kaspar KL, Khoo C, Derrig LH, Schild AL and Gupta K, Consumption of a cranberry juice beverage lowered the number of clinical urinary tract infection episodes in women with a recent history of urinary tract infection. *Am J Clin Nutr* 103:1434–1442 (2016).
- McGuinness SD, Krone R, Metz LM. A double-blind, randomized, placebo-controlled trial of cranberry supplements in multiple sclerosis. *Journal of Neuroscience Nursing.* 2002; 34(1):4-7.
- McKay DL, Blumberg JB. Cranberries (*Vaccinium macrocarpon*) and cardiovascular disease risk factors. *Nutr Rev.* 2007;65:490–502.
- McMurdo ME, Argo I, Phillips G, Daly F, Davey P. Cranberry or trimethoprim for the prevention of recurrent urinary tract infections? A randomized controlled trial in older women. *J Antimicrob Chemother.* 2009; 63(2):389-95.
- McMurdo ME, Bissett LY, Price RJ, Phillips G, Crombie IK. Does ingestion of cranberry juice reduce symptomatic urinary tract infections in older people in hospital? A double-blind, placebo-controlled trial. *Age Ageing.* 2005; 34(3):256-61, <http://dx.doi.org/10.1093/ageing/afi101>.
- Melnyk BM. Integrating levels of evidence into clinical decision making. *Pediatr Nurs* 2004; 30:323–5.
- Mikulic-Petkovsek M, Slatnar A, Stampar F, Veberic R. HPLC-MSn identification and quantification of flavonol glycosides in 28 wild and cultivated berry species. *Food Chem.* 2012;135:2138–46.
- Nova Scotia Department of Agriculture and Fisheries. Cranberry facts and pointers. Accessed online October 14, 2004, at: <http://www.gov.ns.ca/nsaf/elibrary/archive/hort/berrycrops/cranberry/cranfact.htm>.
- Novotny JA, Baer DJ, Khoo C, Gebauer SK and Charrons CS, Cranberry juice consumption lowers markers of cardio metabolic risk, including blood pressure and circulating C-reactive protein, triglyceride, and glucose concentrations in adults. *J Nutr* 145:1185–1193 (2015).
- Pappas, E.; Schaich, K. Phytochemicals of cranberries and cranberry products: Characterization, potential health effects, and processing stability. *Crit. Rev. Food Sci. Nutr.* 2009, 49, 741–781.
- Paquette M, Larque ASM, Weisnagel SJ, Desjardins Y, Marois J, Pilon G et al., Strawberry and cranberry polyphenols improve insulin sensitivity in insulin-resistant, non-diabetic adults: a parallel, double-blind, controlled and randomised clinical trial. *Br J Nutr* 117:519–531 (2017).
- Paredes-López O, Cervantes-Ceja ML, Vigna-Pérez M, Hernández-Pérez T. Berries: improving human health and healthy aging, and promoting quality life—a review. *Plant Foods Hum Nutr.* 2010; 65: 299–308.

- Pérez-López FR, Haya J, Chedraui P. *Vaccinium macrocarpon*: an interesting option for women with recurrent urinary tract infections and other health benefits. *J Obstet Gynaecol Res* 2009;35:630–9.
- Pierre JF, Heneghan AF, Feliciano RP, Shanmuganayagam D, Krueger CG, Reed JD et al., Cranberry proanthocyanidins improve intestinal sIgA during elemental enteral nutrition. *J Parenter Enteral Nutr* 38:107–114 (2014).
- Polashock, J.; Zelzion, E.; Fajardo, D.; Zalapa, J.; Georgi, L.; Bhattacharya, D.; Vorsa, N. The American cranberry: First insights into the whole genome of a species adapted to bog habitat. *BMC Plant Biol.* 2014, 14, 165.
- Prior RL and Gu L, Occurrence and biological significance of proanthocyanidins in the American diet. *Phytochemistry*66:2264–2280 (2005).
- Raz R, Chazan B, Dan M. Cranberry juice and urinary tract infection. *Clin Infect Dis* 2004;38:1413–9.
- Reid G, The role of cranberry and probiotics in intestinal and urogenital tract health. *Crit Rev Food Sci Nutr*42:293–300 (2002).
- Ruel G, Pomerleau S, Couture P, Lamarche B, Couillard C. Changes in plasma antioxidant capacity and oxidized low-density lipoprotein levels in men after short-term cranberry juice consumption. *Metabolism.* 2005;54:856–61.
- Ruel, G.; Pomerleau, S.; Couture, P.; Lemieux, S.; Lamarche, B.; Couillard, C. Favourable impact of low-calorie cranberry juice consumption on plasma HDL-cholesterol concentrations in men. *Br. J. Nutr.* 2006, 96, 357–364.
- Salo J, Uhari M, Helminen M, Korppi M, Nieminen T, Pokka T et al., Cranberry juice for the prevention of recurrences of urinary tract infections in children: a randomized placebo-controlled trial. *Clin Infect Dis*54:340–346 (2012).
- Shmueli H, Yahav J, Samra Z, Chodick G, Koren R, Niv Y, Ofek I. Effect of cranberry juice on eradication of *Helicobacter pylori* in patients treated with antibiotics and a proton pump inhibitor. *Mol Nutr Food Res.* 2007; 51:746–51.
- Stapleton AE, Dziura J, Hooton TM, Cox ME, Yarova-Yarovaya Y, Chen S, Gupta K. Recurrent urinary tract infection and urinary *Escherichia coli* in women ingesting cranberry juice daily: a randomized controlled trial. *Mayo Clin Proc* 2012;87:143–50.
- Stothers L. A randomized trial to evaluate effectiveness and cost effectiveness of naturopathic cranberry products as prophylaxis against urinary tract infection in women. *Can J Urol.* 2002; 9(3):1558-62.
- Sun J, Marais JP, Khoo C, LaPlante K, Vejborg RM, Givskov M et al., Cranberry (*Vaccinium macrocarpon*) oligosaccharides decrease biofilm formation by uropathogenic *Escherichia coli*. *J Funct Foods*17:235–242 (2015).
- Szajdek, A.; Borowska, E. Bioactive compounds and health-promoting properties of berry fruits: A review. *Plant Foods Hum. Nutr.* 2008, 63, 147–156.
- Takahashi S, Hamasuna R, Yasuda M, Arakawa S, Tanaka K, Ishikawa K, Kiyota H, Hayami H, Yamamoto S, Kubo T, et al. A randomized clinical trial to evaluate the preventive effect of cranberry juice (UR65)

- for patients with recurrent urinary tract infection. *J Infect Chemother* 2013;19:112–7.
- Terris MK, Issa MM, Tacker JR. Dietary supplementation with cranberry concentrate tablets may increase the risk of nephrolithiasis. *Urology* 2001; 57:26–9.
- Vasileiou I, Katsargyris A, Theocharis S, Giaginis C. Current clinical status on the preventive effects of cranberry consumption against urinary tract infections. *Nutr Res* 2013; 33:595–607.
- Vicariotto, F. Effectiveness of an association of a cranberry dry extract, D-mannose, and the two microorganisms *Lactobacillus plantarum* LP01 and *Lactobacillus paracasei* LPC09 in women affected by cystitis: A pilot study. *J. Clin. Gastroenterol.* 2014, 48, S96–S101.
- Vinson JA, Bose P, Proch J, Kharrat HA, Samman N. Cranberries and cranberry products: powerful in vitro, ex vivo, and in vivo sources of antioxidants. *J Agric Food Chem.* 2008; 56:5884–91.
- Waites KB, Canupp KC, Armstrong S, DeVivo MJ. Effect of cranberry extract on bacteriuria and pyuria in persons with neurogenic bladder secondary to spinal cord injury. *J Spinal Cord Med.* 2004; 27(1):35–40.
- Wang CH, Fang CC, Chen NC, Liu SS, Yu PH, Wu TY, Chen WT, Lee CC, Chen SC. Cranberry-containing products for prevention of urinary tract infections in susceptible populations. *Arch Intern Med* 2012;172: 988–96.
- Wang CH, Fang CC, Chen NC, Liu SS, Yu PH, Wu TY, Lee CC, Chen SC. Cranberry-containing products for prevention of urinary tract infections in susceptible populations: a systematic review and meta-analysis of randomized controlled trials. *Arch Intern Med.* 2012; 172:988–96.
- Weiss EI, Kozlovsky A, Steinberg D, Lev-Dor R, Bar Ness Greenstein R, Feldman M, Sharon N, Ofek I. A high molecular mass cranberry constituent reduces mutans streptococci level in saliva and inhibits in vitro adhesion to hydroxyapatite. *FEMS Microbiol Lett.* 2004; 232:89–92.
- White BL, Howard LR, Prior RL. Impact of different stages of juice processing on the anthocyanin, flavonol, and procyanidin contents of cranberries. *J Agric Food Chem.* 2011; 59:4692–8.
- Wilson T, Luebke JL, Morcomb EF, Carrell EJ, Leveranz MC, Kobs L, Schmidt TP, Limburg PJ, Vorsa N, Singh AP. Glycemic responses to sweetened dried and raw cranberries in humans with type 2 diabetes. *J Food Sci.* 2010;75:H218–23.
- Wing DA, Rumney PJ, Preslicka CW, Chung JH. Daily cranberry juice for the prevention of asymptomatic bacteriuria in pregnancy: a randomized, controlled pilot study. *J Urol.* 2008; 180(4):1367–72, <http://dx.doi.org/10.1016/j.juro.2008.06.016>.
- Wu X, Beecher GR, Holden JM, Haytowitz DB, Gebhardt SE, Prior RL. Concentrations of anthocyanins in common foods in the United States and estimation of normal consumption. *J Agric Food Chem.* 2006;54:4069–75.
- Xiao SD, Shi T. Is cranberry juice effective in the treatment and prevention of *Helicobacter pylori* infection of mice? *Chin J Dig Dis.* 2003; 4:136–9.

Yung LM, Wong WT, Tian XY, Leung FP, Yung LH, Chen ZY, Yao X, Lau CW, Huang Y. Inhibition of renin-angiotensin system reverses endothelial dysfunction and oxidative stress in estrogen deficient rats. PLoS ONE. 2011;6:e17437.

Zhang L, Ma JL, Pan KF, Go VLW, Chen JS and YouWC, Efficacy of cranberry juice on Helicobacter pylori infection: a double-blind, randomized placebo-controlled trial. Helicobacter 10:139–145 (2005).

ZhuY, Xia M, YangY, LiuF, Li Z, HaoY, Mi M, JinT, Ling W. Purified anthocyanin supplementation improves endothelial function via NO-cGMP activation in hypercholesterolemic individuals. Clin Chem. 2011; 57:1524–33.

