

<https://doi.org/10.46344/JBINO.2022.v11i02.19>

CLINICAL FINDINGS OF PATIENTS WITH HUMAN BRONCHIAL ASTHMA IN BASRAH , IRAQ

Falih Hmood Mezban Ihsan Edan Alsaimary

University of Basrah, College of medicine ,Department of Microbiology, Basrah, Iraq

E-mails: ihsanalsaimary@gmail.com

(Received on date: 17.06.2021

Date of Acceptance: 17.07.2021

Date of publication: 30.02.2022)

ABSTRACT

This study aimed to describe the clinical features of patients asthma in Basrah south in Iraq .The study showed that (3,5) age group population were more affected with asthma (27.9%) and the Females were more affected than males in group 2,3and 5 (6.4%,15.7% and 14.7%)respectively compared to(4.9%,12.3% and 13.2) . in same group of male. While There were (68.6%) of patients came from urban areas in comparison to (31.4%) of cases who came from rural areas. The Smoking patients with positive (43.1%). and well patients with animal contact positive their proportion was while (49%). Seasonal asthma attack in male (23.5%) more than female (20.6%) the perennial asthma attach was recorded in male (29%) more than female (26.9%) in this study show Asthmatic patients with other allergy about (15.7%) and with chronic diseases (31.9%). The percentage of patients with positive family history were 39.2% of the cases, The pulmonary function test result was recorded below (70%) in all age groups. Skin test where the study found highly percentage to HDm to female (76.8%) and (66.2%) to male and HD to female (69.5%) and (68.7%)to male.

Key words: clinical parameters, human bronchial asthma

Introduction

Asthma is a complex respiratory disease in which genetic predisposition, environmental and immunological influences interfere with each other (Edwards ,et al., 2012). It is considered one of the most prevalent chronic diseases, affecting approximately 300 million individuals (Masoli, et al., 2004) and causing an estimated 250,000 deaths each year (Bateman, et al., 2008). In addition, it is projected that by 2025, the global asthma burden will rise by 100 million people due to a growing Westernized lifestyle and urbanization in developing countries (Masoli, et al., 2004). The 'hygiene theory' was originally attributed to an increase in the prevalence of allergic diseases, including asthma, indicating that decreased exposure to microbes during the first years of life plays a role in the development of allergic diseases (Strachan, 1989, 2000). While this theory is generally accepted, studies have shown that the increased incidence of asthma, rhinitis, or Neurodermitis does not completely account for decreased microbial exposure (Mallol, 2008; Brooks ,et al.,2013 and Kramer et al., 2013). Asthma is a widespread illness globally and affects individuals of all ages, This condition usually occurs in infancy and is characterized by variable symptoms of wheeze, dyspnea, and chest tightness caused by air flow obstruction (fully reversible) (GINA, 2015 and Bisgaard & Bonnelykke,2010).

Materials and methods

Samples

A total of (312) patients (149 males and 163 females) of various age groups were included in this Case –control study. The patient was examined, and diagnosed as asthma under supervision of the Physician. The study was carried out during a period from July 2018 to January 2020.

The grouping of patient

Male& Female patients were divided into five groups according to (Falk, 1993; Herd,et al.,1996 ; Nishioka,1996 and charman&Williams,2002)

Group 1: 1- 11 years

Group 2: 12 – 2o years

Group 3: 21- 3o years

Group 4: 31 – 4o years

Group 5: above 4o years

Control group

A total of (204) healthy individual (81 males and 123 females) with out any features of asthma or any allergic to be compared with asthmatic patient in genetic and immunological studies.

Statistical analysis

Statistical analysis is done by using statistical package for social sciences(SPSS) software version 11, the chi square test, univariate and multivariate logistic regression analysis, the ANOVA analysis were applied for correlation between each study parameter, and the difference between two proportion by T-tests were used to assess the significance of difference between groups,P-Value less than 0.05 was considered as Statistically significance(S).P-value < 0.01 as highly significant(HS).and P-value <0.001 as extremely significant(ES).

in 26.9% and 29% other allergies were recorded 6.4% and 9.3% in female and male , family history was recorded 16.6% and 22.5%, and chronic disease were recorded 14,2% and 17.7%. in females and males respectively.

THE RESULTS

Table (1) Show Seasonal asthma attack females and males in percentages 20.6% and 23.5% respectively, while female and male suffering to Perennial asthma attack

Table(1) Clinical characteristic of studied samples.

Clinical characteristic		Female		Male		Total		P.Value	
		NO.	%	NO.	%	NO.	%		
Asthma attack	Seasonal	42	20.6	48	23.5	90	44.1	0.0862	NS
	Perennial	55	26.9	59	29	114	55.9	0.0831	NS
Total		97	47.5	107	52.5	204	100.0	0.0792	NS
Other allergies	No	84	41.1	88	43.2	172	84.3	0.0934	NS
	Yes	13	6.4	19	9.3	32	15.7	0.0732	NS
Total		97	47.5	107	52.5	204	100.0	0.0634	NS
Family history	No	63	30.9	61	30.0	124	60.8	0.0976	NS
	Yes	34	16.6	46	22.5	80	39.2	0.0541	NS
Total		97	47.5	107	52.5	204	100.0	0.0634	NS
Chronic disease	No	68	33.3	71	34.8	139	68.1	0.0734	NS
	Yes	29	14.2	36	17.7	65	31.9	0.0537	NS
Total		97	47.5	107	52.5	204	100.0	0.0634	NS

Discussion

Asthma attacks, according to the study majority (55.9%) in perennial were (44.1%) in seasonal, the researcher believe this result due to increasing environmental pollution in addition to lack of concern for health status. These result confirmed by **Tajiri,(2014)**.

The study found that the number of asthma patients with a family history was 39.2%. (Table 1). Cengizlier and Misirlioglu posted similar findings (2006).All of the recent asthma family history studies confirmed that asthma was strongly associated with family history of asthma. **Fuhlbrigge, (2001)** mentioned that about three-fourths of all cases of asthma seem to be hereditary. **Weiss, (2001)** reported that risk of asthma increases in the children if one of the parent being asthmatic also it is more likely if both parents are asthmatic. Even identical twins are more likely being asthmatic than non-identical twins. **(Mingomataj et al., 2008)** found that the percentage of family history was 23.3%. **(Eseverri et al., 1998)** found out that among the risk factors of asthma, 69% of the patients had a family history.

With respect to family history (39.2%) of the subjects had the family history of asthma. The finding of this study indicates that less than half of the studied sample has a healthy bronchial asthma family history, the researcher suggests that this outcome may suggest that heredity plays a major role in disease. This finding was in line with the findings of several studies that showed that bronchial asthma is a family disease and can also be a script for

external aggravating factors. Backed by the results of **(Parvin, et al., 2011)**. This finding and research showed that most of the studied individuals had a family history of asthma (49.3 percent).

The analysis revealed a high percentage (84.3 percent) of the study samples of other allergies not present (15.7 percent) from samples of other allergies present because the researcher assumes this finding is attributable to experience of asthma patients who are subjected to continuous weekly

Treatment at the Basra Allergy and Asthma Center. **(Hong,et al, 2012)**. did not confirm these outcomes , The majority (68.1 percent) of the study was not present with respect to chronic illness, although (31.9 percent) were present, the researcher assumes that this outcome is due to increased samples taken from women in age groups 3 and 5 and the hormone variations between the sexes. Such outcomes are confirmed by From **Nafstad, (2001)**.

These results of present study were not confirmed by with the finding of many studies done by **(Ronmark, et al., 2005; Nicolai, et al., 2001 ; Becklake, and Kauffmann, 1999)**. Another study **(Anthony , et al., 2004)** was entitled with “ New-onset asthma among soldiers serving in Iraq and Afghanistan “ whereas their results show that the women had higher risks of asthma: 1.90 (1.37, 2.63) for men, and 1.70 (0.66, 4.40) for women.

The study found that the number of asthma patients with a family history was 39.2%. (table 1). **Cengizlier and Misirlioglu** posted similar findings **(2006)**. All of the recent asthma family history studies confirmed that asthma was strongly associated with family history of asthma. **Fuhlbrigge, (2001)** mentioned that about three-fourths of all cases of asthma seem to be hereditary. **Weiss, (2001)** reported that risk of asthma increases in the children if one of the parent being asthmatic also it is more likely if both parents are asthmatic. Even identical twins are more likely being asthmatic than non-identical twins . **Mingomataj et al., (2008)** found that the percentage of family history was 23.3%. **Eseverri et al., (1998)** found out that among the risk factors of asthma, 69% of the patients had a family history.

With respect to family history (39.2%) of the subjects had the family history of asthma. The finding of this study indicates that less than half of the studied sample has a healthy bronchial asthma family history, the researcher suggests that this outcome may suggest that heredity plays a major role in disease. This finding was in line with the findings of several studies that showed that bronchial asthma is a family disease and can also be a script for external aggravating factors. Backed by the results of **Parvin, et al., this finding (2011)**. The research showed that most of the studied individuals had a family history of asthma (49.3 percent).

The analysis revealed a high percentage (84.3 percent) of the study samples of other allergies not present (15.7 percent)

from samples of other allergies present because the researcher assumes this finding is attributable to experience of asthma patients who are subjected to continuous weekly

Treatment at the Basra Allergy and Asthma Center . **Hong,et al**, did not confirm these outcomes **(2012)**.

The majority (68.1 percent) of the study was not present with respect to chronic illness, although (31.9 percent) were present, the researcher assumes that this outcome is due to increased samples taken from women in age groups 3 and 5 and the hormone variations between the sexes. Such outcomes are confirmed by **From Nafstad, P (2001)**. These results of present study were not confirmed by with the finding of many studies done by **(Ronmark,et al., 2005 ; Nicolai, et al., 2001 ; Becklake, and Kauffmann, 1999)**, Another study **Anthony , et al., (2004)** was entitled with " New-onset asthma among soldiers serving in Iraq and Afghanistan " whereas their results show that the women had higher risks of asthma: 1.90 (1.37, 2.63) for men, and 1.70 (0.66, 4.40) for women.

Reference

- Aaron, S. D.; Vandemheen, K. L.; FitzGerald, J. M.; Ainslie, M.; Gupta, S.; Lemièrre, C.; Field, S. K.; McIvor, R. A.; Hernandez, P. and Mayers, I. (2017)**. Reevaluation of diagnosis in adults with physician-diagnosed asthma. *JAMA.*, 317(3), 269–279.
- Adams, K. E. and Rans, T. S. (2013)**. Adverse reactions to alcohol and alcoholic beverages.

Annals of Allergy, Asthma and Immunology, 111(6), 439–445.

Agache, I., Ciobanu, C., Agache, C. and Anghel, M. (2010). Increased serum IL-17 is an independent risk factor for severe asthma. *Respiratory medicine*, 104(8):1131-1137.

Agarwal, R. (2011). Severe asthma with fungal sensitization. *Current Allergy and Asthma Reports*, 11(5), 403.

Ahmadi, M.; Rahbarghazi, R.; Aslani, M. R.; Shahbazfar, A. A.; Kazemi, M. and Keyhanmanesh, R. (2017). Bone marrow mesenchymal stem cells and their conditioned media could potentially ameliorate ovalbumin-induced asthmatic changes. *Biomedicine & Pharmacotherapy*; 85: 28–40.

Al Niami, B.F.(1990) . immunological assessment of adult male asthmatic patients. MSc Dissertation, University of Baghdad .

Alnahas, S.; Hagner, S.; Raifer, H.; Kilic, A.; Gasteiger, G.; Mutters, R.; Hellhund, A.; Prinz, I.; Pinkenburg, O.; Visekruna, A. and Garn, H. (2017). IL-17 and TNF- α are key mediators of *Moraxella catarrhalis* triggered exacerbation of allergic airway inflammation. *Frontiers in immunology*. 8: 1562.

Alsaimary, I.(2006). Atopic dermatitis/ Eczema Syndrome : A Molecular investigation of microbial super antigenicity with bacteriological , immunological ,clinical ,hematological and histopathological studies.Basra University,college of medicin

Alyasin, S., Karimi, M.H., Amin, R., Babaei, M. and Darougar, S.(2013). Interleukin-17 gene expression and serum levels in children with severe asthma. *Iranian journal of immunology*, 10(3):177-185.

Amelink, M.; De Nijs, S.B.; De Groot, J.C.; Van Tilburg, P.M.; Van Spiegel, P.I.; Krouwels, F.H.;

Lutter, R.; Zwinderman, A.H.; Weersink, E.J.; Ten Brinke, A.; Sterk, P.J. and Bel, E.H.(2013). Three phenotypes of adult-onset asthma *Allergy*: 68(5): 674-680.

Barczyk, A.; Pierzchala, W.; Sozanska, E.(2003). Interleukin-17 in sputum correlates with airway hyperresponsiveness to methacholine.*Resp. Med*, 97:726-733.

Barnes, P. J. (2008). The cytokine network in asthma and chronic obstructive pulmonary disease. *The Journal of Clinical Investigation*, 118(11), 3546–3556.

Barnett, S.B. and Nurmagambetov, T.A.(2011). Costs of asthma in the United States: 2002-2007. *J. Allergy Clin. Immunol.*;127(1):145-52.

Bateman, E. ; Hurd, S.; Barnes, P.; Bousquet, J.; Drazen, J. ; FitzGerald, M.; Gibson, P.; Ohta, K.; O'byrne, P. and Pedersen, S E.,(2008). Global strategy for asthma management and prevention: GINA executive summary. *Eur. Respir. J.*;31(1):143-78.

Bateman, E. D.; Hurd, S. S.; Barnes, P. J.; Bousquet, J.; Drazen, J. M.; FitzGerald, M.; Gibson, P.; Ohta, K.; O'byrne, P. and Pedersen, S E.(2008).The Global strategy for asthma Management and prevention:GINA Executive summary. *Eur. Res. J.* ;31(1):143-78

Batra, V.; Musani, A. I.; Hastie, A. T.; Khurana, S.; Carpenter, K. A.; Zangrilli, J. G. and Peters, S. P. (2004). Bronchoalveolar lavage fluid concentrations of transforming growth factor (TGF)- β 1, TGF- β 2, interleukin (IL)-4 and IL-13 after segmental allergen challenge and their effects on α -smooth muscle actin and collagen III synthesis by primary human lung fibroblasts. *Clinical and Experimental Allergy*, 34(3), 437–444.

Mezban .Falih hmood and alsaimary Ihsan edan . 2016 . Significance of Skin test reactivity

to aeroallergens in patients with chest and skin allergies in basrah through 2014 and 2015. Basic Research Journal of Microbiology. Vol. 3(6)

