

<https://doi.org/10.46344/JBINO.2021.v10i04.16>

DEPRESSION AND ANXIETY FREQUENCY IN PATIENTS HOSPITALIZED ON THE GUADALAJARA REGIONAL MILITARY HOSPITAL IN THE MONTH OF APRIL 2019.

Miranda Nava, Gabriel¹. MD. Gallo Frías, Luis Gilberto². MD.

1. Major of military armed forced of Mexico. Neurologist.

2. Medical Doctor. Social Service. Lamar University. Guadalajara.

Email: drgabrielmiranda@hotmail.com

ABSTRACT

Observe and Identify patients that presented depression and anxiety using the Hospital Anxiety Diagnosis Scale (HADS), Zung & Conde scale and ASQ 15 scale on Guadalajara Regional Military Hospital during the month of April 01st to April 30th 2019. Methods: This is a cohort, non experimental, observational, prospective and longitudinal study with PubMed and NCBI articles as variables. Findings: Most patients presented anxiety, most patients had a chronic illness, depression was seen mostly in patients older than 50 years old, there was inadequate diet, lack of sleep, and low distress level.

Keywords: Nil

INTRODUCTION

Depression and anxiety affect most people around the world, it is characterized by a presence of fear, loss of interest, feelings of guilt or self-esteem that are more commonly associated with sleep disorders, lack of appetite, lack of energy or difficulty concentrating. Depression can become chronic or recurrent and difficult the overall performance on a daily basis, or capacity to live day by day, in its most dangerous form it can lead often to suicide and its lowest form it can be treated with medication and professional psychotherapy. (salud, Depression , 2017)

Anxiety is one of the major disorders and its characterized by persistent concern during any activity or routine it is difficult to treat and it can affect the way a person feels physically. (mayoclinic, 2018)

During this investigation we will observe a sample that was taken on the Guadalajara Regional Military Hospital during the month of april a sample of 56 patients presented anxiety and depresión according to three scales that were applied.

Hads (Hospital Anxiety Diagnosis Scale)

The Hospital Anxiety Diagnosis Scale is an autoapplicable questionnaire integrated by 14 items with subscales of seven items one for impaired questions and one with pair questions for depresión, the authors for this scale are Zigmund and Snaith who proposed this in 1983 and defined the concepts of anxiety and depresión the objective of this scale is to identify if the patient has being tensed, concerned or frightened in any way, the 8 items that form the depression subscale are centered around anhedonia with a maximum score that binds from 0 to a 39 score, in which 0-9

score means lack of stress, 10-19 means low stress, 19 to 29 means mild stress and 30 to 39 means anxiety and severe depression.

Zung & Conde Scale

Its and autoapplicable scale consisting of 20 phrases related to depression formed by 10 negative phrases and 10 positive phrases which relate to strong somatic symptoms and 8 cognitive items for each group contemplating the scale with two items refere to mood and other psychotic symptoms. (Roza, 2019)

Depression and Anxiety

Severe Depression: Its characterized by a combination of symptoms that interfere with capacity to work, sleep, study, eat and enjoy daily basis activities.

Dysthimic Disorder

Its characterized by simptoms that is somewhat between 2 years and beyond but less severe, it incapacitates the patient and it prevents him from having a normal life accompanied by a severe depression episode during life.

Psychotic depression

Occurs during severe depression and its accompanied by some for of psicosis accompanied by delirium and allucinations.

Seasonal Depression

Its characterized by depresión that appears during Winter or times of decreased sunlight.

Bipolar Disorder

Its characterized by maniac depresión disorder that its accompanied by cyclic mood swings and depression state, its often seen in patients with cáncer, HIV/Aids and Parkinson.

SYMPTOMS

Emotional

Are accompanied by guilt ideas, a severe disease, ideas of sadness never going to heal, loneliness, lack of concentration because patient will eventually die.

Physical

Difficulty eating, or basic needs, weight loss, mood swings.

Negative thoughts

This is mostly seen in older patients, self-esteem problems, most cases are seen in patients over 60 years old, or below 45 years old.

METHODS

This is a cohort, non experimental, observational, prospective and longitudinal study in which scholarship was evaluated, cause of hospitalization, age, previous diseases, job and the days patient had being hospitalized.

56 patients both men and women older than 18 years old were evaluated during this study, a random sample was taken in which every patient has the same possibility of presenting depression or anxiety. Patients hospitalized in the womens hospital room, the mens hospital

room, and the room that consisted of patients that had the rank of major in the mexican armed forces or above excluding patients that belonged to Intensive care unit, using the Hospital Anxiety Diagnosis Scale, Zung & Conde Scale and ASQ-15 Scale were used during this study.

JUSTIFICATION

This study was conducted to observe what was the impact of being hospitalized and the relation it had with depression and anxiety in patient, we pretended to find viable date that allowed us to expose the hospital enviroment and the presence of disease, anxiety and depression.

RESULTS

Figure 1. Patients with depression. Fuente. Zung & Conde Depression Scale. 62% of the patients did not present depression 34 patients, 16 patients had low depression 29%, also 8% of the patients had moderate depression which represented 4 patients also 2 of the patients representing 1% presented what could be considered as severe depression.





Figure 2. Most common ages seen during study.



42% of the patients presented depression were over 50 years old.

35% of the patients presented depression between 40-50 years of age.

17% of the patients that presented depression had between 30-40 years of age.

6% of the patients had between the ages of 20 to 30 years of age.




Figure 3. Patients that had some sort of stress evaluated by HADS scale.

43% of the patients had lack of stress.

46% of the patients had low level stress.

11% of the patients had moderate stress.

0% of the patients had severe stress.




Figure 4 Most common diseases (12 patients didnt have an illness):

- Hipertensión: 16
- Diabetes Mellitus:12
- Renal Insuficiency: 4
- Ulcerative Colitis:1
- Hepatic Cirrosis: 3
- Lung Cancer: 3
- Cervical Cancer:1
- Fractures: 4




Figure 5. Civil Status:

- Married: 54%
- Separated: 35%
- Single: 11%




Figure 6. Patients that presented anxiety

- 64% presented anxiety 35 patients
- 36% did not present anxiety 21 patients




Figure 6. Balanced Diet:

- 86% No
- 14% Yes

CONCLUSION

Referred to this subject we understand this two disorders are preventable and the patient if its treated and seeks help with time he can change his daily habits, our suggestions are that a stable lifestyle with a well balanced diet consisting of fruit, vegetables, meat, daily exercise, stable relationships with family and friends, alongside no work stress, in addition to a good mental stability can lead to a good life and to prevent this type of disorders

also to prevent chronic illness which were seen during this study on most patients that indicated feeling anxious or depressed, if they are in this state also to take medication on time and with the help of family members and friend.

- We observe most patients presented a low depression level
- Most patients presented anxiety
- Most patients had a chronic illness
- Patients over 50 years old presented higher depression levels

- Most patients were married
- Most patients had low level stress level.

http://ipsi.uprrp.edu/opp/pdf/materiales/aniedad_esp.pdf

REFERENCES

- 1) Carolina Casanova. (2017) . Screening for symptoms of anxiety and depression in patients admitted to a university hospital with acute coronary syndrome.Revisado en abril de 2019. http://www.scielo.br/scielo.php?script=sci_abstract&pid=S2237-60892017000100012&lng=en&nrm=iso
- 2) Grupo de Trabajo de Guia de Practica Clinica para el manejo de pacientes con trastornos de ansiedad en atención primaria. Plan Nacional para SNS. (2008) Guías de practica clínica en SNS http://www.guiasalud.es/GPC/GPC_430_Ansiedad_Lain_Entr_compl.pdf
- 3) Servicio de Psiquiatria y Psicología Hospital Universitario Dexeus (2016) Barcelona. <http://www.psiquiatriapsicologia-dexeus.com/es/unidades.cfm/ID/1099/ESP/tipos-depresion.htm>
- 4) National Institute of Mental Health. Departamento de Salud y Servicios Humanos de los Estados Unidos. Trastornos de Ansiedad.
- 5) Enfocando la depresión como problema de salud publica en México (2012) Fernando A. Wagner. Catalina González http://www.scielo.org.mx/scielo.php?script=sci_arttext&pid=S0185-33252012000100002
- 6) Guillermo Hernandez. Grisel. Monica. Kimmelman. (2005) Orellana Rev. méd. Chile v.133 n.8 Santiago ago. 2005 Trastornos de ansiedad en pacientes hospitalizados en Medicina Interna https://scielo.conicyt.cl/scielo.php?script=sci_arttext&pid=S0034-98872005000800005
- 7) Miriam Costas. Veronica Prado. Jose Manuel Crespo. (2013). Ansiedad y depresión entre los pacientes hospitalizados en el complejo hospitalario de Ferrol. <file:///C:/Users/user/Downloads/Dialnet-AnsiedadYDepresionEntreLosPacientesHospitalizadosE-4394399.pdf>
- 8) Sílvia Abduch Haas^{1,*}; Daniela Centenaro Levandowski² ; Antônio Nocchi Kalil. (2017). Ansiedad, depresión, percepción y sentimientos de mujeres con cáncer indicadas para cirugía de exenteración pélvica

<file:///C:/Users/user/Downloads/57086-Texto%20del%20art%C3%ADculo-114521-2-10-20171011.pdf>

- 9) YeLi.Mei-RongLv^bYan-JinWei^cLingSun^bJi-XiangZhang^dHuai-GuoZhang^e (Julio de 2017) Dietary patterns and depression risk: A meta-analysis <https://www.sciencedirect.com/science/article/abs/pii/S0165178117301981>
- 10) General IMSS. Consejo de Salubridad Guia de referencia rapida guia practica clínica http://sgm.issste.gob.mx/medica/medicadocumentacion/guiasautorizadas/Psiquiatria/IMSS-161_09_TRASTORNO_DEPRESIVO/GRR%20Trastorno%20Depresivo.pdf
- 11) Organización Mundial de la Salud. Que es depresión y ansiedad. www.who.com
- 12) Roza. M. (2019) escala autoadministrada de depresion psicomag. Escala de Zung y Conde que es. <http://www.depresion.psicomag.com/esczung.php>