

STUDY OF ANTIMICROBIAL ACTIVITY *AGERATUM CONYZOIDES* & *OCIMUM GRATISSIMUM*

Mohammad Chand Jamali

Department of Health & Medical Sciences, Khwarizmi International College, 25669, Abu Dhabi, United Arab Emirates,

ABSTRACT

Antioxidants are both natural and synthetic compounds with the ability to scavenge free radicals and inhibit oxidation processes [3]. Many synthetic compounds have ability to scavenge free radicals (such as Butylated Hydroxy Anisole (BHA), and Butylated Hydroxy Toluene (BHT)) are very effective and are used for industrial processing but they possess some side effects and toxic properties to human health. Anti-oxidants play an important function in protecting the body against free radicals. They achieve this by stopping the formation of new free radicals species. In the present experiment we are studying the antimicrobial activity of *Ageratum Conyzoides* and *Ocimum Gratissimum*.

Keywords: Anti Microbial, *Ocimum Gratissimum*, *Ageratum Conyzoides*



INTRODUCTION

There are several diseases of the gastrointestinal tract of farm animals in which diarrhea is a major clinical finding. Diarrhea is the increased frequency of defecation of watery faeces. Commonly incriminated bacterial agents of diarrhea include *Escherichia coli*, *Salmonella* spp, *Clostridium perfringens* types B & C, and *Mycobacterium paratuberculosis*. Bacterial agents induce diarrhea in newborn animals particularly calves and piglets in the first week of life. It is one of the most common disease complexes which the large animal clinicians have to contend with. It causes a significant economic loss in herds of cattle and swine. It may assume even greater importance in the future as livestock production becomes more intensified. *Escherichia coli* serovars are the causative agents of colibacillosis. The utilization of antimicrobial drugs has played an important role in animal husbandry, since they are used in prophylaxis, treatment and growth promotion. In the developed world, the extensive use of antibiotics in agriculture, especially for prophylactic and growth promoting purposes, has generated much debate as to whether this practice contributes significantly to increased frequencies and dissemination of resistance genes into other ecosystems. In developing countries like Nigeria, antibiotics are used only when necessary, especially if the animals fall sick, and only the sick ones are treated in such cases. There is currently a world trend to

reduce the use of antibiotics in animal food due to the contamination of meat products with antibiotic residues, as well as the concern that some therapeutic treatments for human diseases might be jeopardized due to the appearance of resistant bacteria. During the last decade, the use of antimicrobial drugs for growth promotion and therapeutic treatment in food animals has received much attention. The effectiveness of currently available antibiotics is decreasing due to the increasing number of resistant strains causing infections. The reservoir of resistant bacteria in food animals implies a potential risk for transfer of resistant bacteria, or resistance genes, from food animals to humans. Subsequent emergence of infections in humans, caused by resistant bacteria originating from the animal reservoir, is of great concern. These unintended consequences of antimicrobial drug use in animals led to termination of antimicrobial growth promoters in food animals in countries in the European Union, including Denmark, where the consumption of antimicrobial drugs by production animals was reduced by 50% from 1994 to 1997. The effective treatment and control of diarrhea in cattle and piglets can be unreliable. Considerable progress had been made in the treatment of the effects of diarrhea, such as dehydration and acidosis but less so in the control of these disease complexes.

There have been several approaches to combat diarrhea in livestock among these are the resurgence of interest in the use of medicinal plants and this subject has received extensive coverage in many publications. This has a wide range of benefits to man. From trees and shrubs amongst others, man obtains medicine and foods. A good number of trees and shrubs have been claimed by ethnoveterinary practitioners, ethnomedical practitioners and other local people to have medicinal benefits against infectious and/or non infectious animal and/or human diseases. Throughout history plants have been used by human beings for medicinal purposes and even in modern times have formed the basis of many pharmaceuticals in use.

A recent review has shown that approximately 25% of modern medications are plant derivatives, while 75% of new drugs against infectious diseases that have arrived between 1981 and 1997 originated from natural sources. Medicinal plants are those plant from which one or more of its tissues contain substances that can be used for the therapeutic purposes or which are precursors for the synthesis of useful drugs. It is possible to distinguish between medicinal plant whose constituents have well established medicinal properties and plants whose constituents have not been thoroughly evaluated for therapeutic purposes.

Despite the progresses in modern medicine, it has been reported that more than 70% of the developing world's population still depends on

complementary and alternative systems of medicine, otherwise known as traditional medicine. Many of these medicines are plant based, and include curcumin, resveratrol, baicalein, boswellic acid, betulinic acid, ursolic acid, and oleanolic acid are now studied as possible drugs for the future against inflammatory diseases. Edible plant extracts have shown promising anti-tumorigenic activity. A large number of traditional medicinal preparations used in various countries of the world have shown promise as treatment for herpes virus infections, which cannot be completely cured by the available antiherpes drugs like nucleoside analogs. The various ailments for which searches are going on in plants for newer drugs are too numerous to mention; to cite just a few examples, new plant-derived compounds are being searched for treatment of malaria because of the increasing resistance of *Plasmodium falciparum* against currently available anti-malarial drugs, searches for anti-fungal, molluscicidal and larvicidal compounds in African medicinal plants have been reviewed, anti-leishmanial activity in Israeli plants has been reviewed by El-On et al., and anti-trypanosomal and cytotoxic activities of pyrrolizidine alkaloid-producing plants of Ethiopia has been reported by Nibret et al. The various ailments treated by the plants included respiratory tract infections, gastrointestinal disorders, sprains and fractures, skin ailments, malaria, mental disorders, hepatic disorders, diabetes, influenza, urinary tract infections, hypertension, debility, measles, chicken pox, toothache, gynecological problems, sexual disorders, and helminthiasis. Some

plants were also used as antidote to poison, diuretic, and abortifacient.

Efforts are now being made in different countries to carry out research into traditional medicine and to integrate it with orthodox medicine. However, not much has been done to salvage traditional medicine as it applies to veterinary practice. Although, there are no professional traditional veterinary medical practitioners in Nigeria, herd men, village elders and people who kept animals for domestic purposes were experienced in the diagnosis and treatment of specific animal diseases. Some of these herbs are used mainly for the treatment of diseases found in animal only. Nigerians have developed a stock of empirical information concerning the therapeutic values of our local plants, and it is widely believed that remedies of natural origin are harmless, and carry no risk. However, there is a possibility of inadvertently applying extracts of toxic plants in a few cases.

It is therefore important that the

MATERIALS AND METHODS

Collection of samples

The various medicinal plants were obtained at the campus of the University of Ibadan. They include: *Ageratum conyzoides* *LOcimum gratissimum* L.

Extraction of the medicinal plants

Two (2) extracts were obtained from each of the medicinal plants namely; chloroform, methanol and water extracts. Chloroform and methanol base extraction involved three major processes, using the

knowledge of traditional treatments of animal diseases be documented and detailed studies be carried out to establish the efficacy of these herbs. Also many of the medicinal plants acclaimed to be of therapeutic value are yet to be completely exploited hence a need for further investigation in order to determine the chemical composition and the efficacy of extracts of the plants in combating diseases. Phytochemical studies have shown that plants with antimicrobial activity contain bioactive constituents such as tannins, flavonoids, alkaloids and saponins.

The study aimed at isolating and identifying *Escherichia coli* from diarrhoeic fecal samples of calves and piglets in Ibadan, Nigeria. It also aimed at extracting the therapeutic component of the medicinal plants using different solvents and determining the efficacy of the plants extracts on *E.coli* by disc diffusion and minimal inhibitory concentration (MIC) methods.

continuous alcohol extraction method with slight modification. The plants were subjected first to chloroform base extraction and then to methanol base extraction.

Standardization of E. coli Used for Disc

Diffusion Susceptibility Test

A discrete colony was inoculated into tryptone soya broth and incubated aerobically at 37°C for 18 to 20hrs. The broth was diluted by adding a loopful of the overnight culture to 5.0ml of physiological saline contained in bijou

bottle. This gave a concentration of approximately 10^5 - 10^6 CFU per ml. With a sterile Pasteur pipette, about 2.0ml of the diluted suspension was transferred to sensitivity test Agar. The plate was hand rocked back and forth and side ways to allow the culture to spread over the whole surface. The excess fluid was removed with a Pasteur pipette and discarded into a strong disinfectant. The plate was left to dry in an inverted position on the working bench.

Determination of the effects of the Extracts

Discs of about 5.0mm in diameter were prepared from Whatmann filter topic. About 0.05ml of each of the plant extracts was then impregnated into the disc by a single channel micro pipette. The discs were then applied onto the surface of the already inoculated agar. The plate was left on the working bench to allow for the diffusion of the plant extract into the agar before incubating at 37°C for 20hrs. The test was carried out for chloroform, methanol and water extracts of the plants. Zones of inhibition was then observed and measured in millimeter after aerobic incubation at 37°C for 18 – 20hrs.

Determination of the MIC of the Plant Extract on E. coli Using Tube Dilution Method

The standard inoculums used were determined by Miles and Mistra method. The minimum inhibitory concentration (MIC) test was carried out only for *Ocimum gratissimum*. There E. coli strains were used for the test. The strains were designated 'A', '4' and '5'. 'A' was isolated from calf

faecal sample, '4' is E. coli 021, strain, while, sample '5' is E. coli 022 strain. Trypticase soy broth was delivered into a row of 9 test tubes for each of the strains. About 0.9ml of the broth was delivered into the first test tubes and about 0.5ml of the broth was delivered into the rest of the test tubes. Also, 0.1ml of known concentration of methanol extract of *Ocimum gratissimum* was then delivered into the first test tube. The concentration of the *Ocimum gratissimum* used was $2.0 \times 10^4 \mu\text{g/ml}$. A doubling dilution was then carried out down the row of the test tubes except the 9th. This was done by transferring 0.5ml of the mixture in the first tube into the second tube. This dilution was repeated in the third test tube as well as in the other tubes until the last was reached. The 0.5ml of the mixture in 8th tube was discarded. The 9th test tube which served as a control contained 0.5ml of broth without the extract. This procedure gave antibiotic concentration of $2.0 \times 10^4 \mu\text{g/ml}$ in the first test, $1.0 \times 10^4 \mu\text{g/ml}$ in the second and so forth. Equal volume, that is, 0.4ml of the prepared diluents of the organism was then added to the row of the test tubes. The preparation was incubated at 37°C for 24hrs after which the content of the tubes were inoculated onto blood agar and then incubated aerobically at 37°C for 24 hrs.

Chloroform extract, ethanol extract, and water extract of all the medicinal plants were tested on three (3) strains of the E. coli. Growth was determined by visual observation for turbidity and by planting a loopful of the broth containing the E. coli and the extract of *Ocimum*

gratissimum on blood Agar and incubated aerobically at 37°C for 24hrs.

RESULTS

Susceptibility of E. coli Strains to Plants Extracts by Diffusion Test

Two of the plants namely Ageratum conyzoides and Ocimum gratissimum however were observed to show some degree of inhibition which varies with the E. coli strain tested. The results of the

susceptibility of E. coli strains to plants extracts are as shown in Table 1 The activities of the extracts also varied between solvents with the ethanol extracts demonstrating the highest activity against all the test bacteria.

The methanol extracts of the plant parts were more potent than the chloroform and water extracts which showed no activity against the test organisms (Table 2).

Table 1: Antimicrobial effects of plant extracts on E. coli strains

Plants	Strain of E. coli	Chloroform	Methanol	Water
Ageratum conyzoides	A	-	10mm	-
	4	-	-	-
	5	-	12mm	-
Ocimum gratissimum	A	-	13mm	-
	4	-	12mm	-
	5	-	13mm	-

Legend: A = isolated strain from diarrhoeic pig; 4 = E. coli 021; 5 = E. coli 022; - = no inhibition

Table 1 shows the MIC of the extract of Ocimum gratissimum on E. coli strain A. It showed that the extract of Ocimum gratissimum inhibited E. coli strain A at MIC value of 20,000µg/ml.

Minimum Inhibitory Concentration (MIC) Test

Table 2 shows the MIC of the extract of *Ocimum gratissimum* on *E. coli* strain 4. It showed that the extract of *Ocimum gratissimum* also inhibited *E. coli* strain 4 at MIC value of 20,000 µg/ml.

Table 3 shows the MIC of the the extract of *Ocimum gratissimum* on *E. coli* strain 5. This also showed that the extract of *Ocimum gratissimum* on inhibited *E. coli* strain 5 at MIC value of 20,000 µg/ml.

Table 2: Minimum Inhibitory Concentration of *Ocimum gratissimum* on *E. coli* Strain A

E. coli strain A load (Cfu/ml)	Concentration of <i>Ocimum gratissimum</i> µg/ml	Growth Observed
7.75 x 10 ³	20,000	No growth
7.75 x 10 ³	10,000	Presence of growth
7.75 x 10 ³	5,000	Presence of growth
7.75 x 10 ³	2,500	Presence of growth
7.75 x 10 ³	1,250	Presence of growth
7.75 x 10 ³	625	Presence of growth
7.75 x 10 ³	312.5	Presence of growth
7.75 x 10 ³	156.25	Presence of growth
7.75 x 10 ³	0.0	Presence of growth

Table 3: Minimum Inhibitory Concentration of *Ocimum gratissimum* on *E. coli* Strain 4

E. coli strain 4 load (Cfu/ml)	Concentration of Ocimum gratissimum µg/ml	Growth Observed
1.13 x 10 ⁴	20,000	No growth
1.13 x 10 ⁴	10,000	Presence of growth
1.13 x 10 ⁴	5,000	Presence of growth
1.13 x 10 ⁴	2,500	Presence of growth
1.13 x 10 ⁴	1,250	Presence of growth
1.13 x 10 ⁴	625	Presence of growth
1.13 x 10 ⁴	312.5	Presence of growth
1.13 x 10 ⁴	156.25	Presence of growth
1.13 x 10 ⁴	0.0	Presence of growth

Table 4: Minimum Inhibitory Concentration of *Ocimum gratissimum* on *E. coli* Strain 5

E. coli strain 5 load(Cfu/ml)	Concentration of Ocimum gratissimum µg/ml	Growth Observed
1.18 x 10 ⁴	20,000	No growth
1.18 x 10 ⁴	10,000	Presence of growth
1.18 x 10 ⁴	5,000	Presence of growth
1.18 x 10 ⁴	2,500	Presence of growth
1.18 x 10 ⁴	1,250	Presence of growth
1.18 x 10 ⁴	625	Presence of growth
1.18 x 10 ⁴	312.5	Presence of growth
1.18 x 10 ⁴	156.25	Presence of growth
1.18 x 10 ⁴	0.0	Presence of growth

DICUSSION

In the tropical environment, diarrhea is responsible for mortality in young animals especially piglets. Survivors lose weight and may not be able to reach weaning weight due to atrophy of the intestinal mucosa. The Enteropathogenic *E. coli* remains an important cause of neonatal diarrhea. This has been confirmed by the findings of other workers. *E. coli* coliforms make-up approximately 10% of microorganism of humans and is used as indicator organisms of faecal contamination and is associated with poor environmental sanitation. However, many drug-resistant human fecal *E. coli* isolates may originate from poultry, whereas drug-resistant poultry-source *E. coli* isolates likely originate from susceptible poultry-source precursors. It is estimated that nearly 90% of all antibiotic agents use in food animals, are given at subtherapeutic concentrations prophylactically or to promote growth. Antimicrobial drug resistance in *E. coli* isolated from wild birds has been described. *E. coli* and the *E. coli* 0157: H7 strain has previously been isolated from meat samples.

E. coli strain showed a high frequency of single or multiple drug resistance (MDR). Multidrug resistance was defined as resistance to ≥ 3 of the antimicrobial agents tested, which revealed the multi-drug resistant pattern and ability of these organisms to many antimicrobials (3 to 6 antibiotics). The bacteria were found to be resistant to Ampicillin, Nalidixic acid, Nitrofurantoin, Co-trimoxazole, Streptomycin and

Tetracycline. The MDR pattern reported on *E. coli* in this study is comparable to previous studies. One suspected source of drug-resistant *E. coli* in humans is use of antimicrobial drugs in agriculture. This use presumably selects for drug-resistant *E. coli*, which may be transmitted to humans through food supply. Supporting this hypothesis is the high prevalence of antimicrobial drug-resistant *E. coli* in retail meat products, especially poultry reported by Schroeder et al. and Johnson et al., and the similar molecular characteristics of fluoroquinolone-resistant *E. coli* from chicken carcasses and from colonized and infected persons in Barcelona, Spain, in contrast to the marked differences between drug-susceptible and drug-resistant source isolates from humans. Such multidrug resistance has important implications for the empiric therapy of infections caused by *E. coli* and for the possible co-selection of antimicrobial resistance mediated by multidrug resistance plasmids.

E. coli contamination is usually associated with contaminated water and food (animal feeds) and their presence reflects the degree of purity or signals faecal contamination of both human and animal origin. The presence of *E. coli* in these diarrheic piglets should receive particular attention, because their presence indicate public health hazard and give warning signal for the possible occurrence of food borne intoxication. Although the number of isolates was relatively small, the level of multidrug resistance was high. The high rates of resistance found in the present study can

be explained by the spread of use of antibiotics agents given to farm animals in Nigeria as prophylaxis, growth promoters or treatment. According to Abdellah et al and Okonko et al., the multiple resistances observed were also to those antimicrobials frequently employed in veterinary practices. Therefore, in line with the assertions of Okonko et al., we recommend that more restrictions on the irrational use of antibiotics and public awareness activities should be undertaken to alert the public of the risks of the unnecessary use of antibiotics.

In a study by El-Mahmood and Doughari, *S. aureus*, *S. pyogenes* and *P. mirabilis* were more susceptible to *Cassia alata*, while *E. coli* and *P. aeruginosa* were less sensitive. Two of the plants (*Ageratum conyzoides* and *Ocimum gratissimum*) showed some degree of inhibitions with ethanol extract of the plant. This is consistent with previous studies by various authors. Paste of the whole plant of *Ageratum conyzoides* is used as an insect repellent and for treatment of wounds and itches. Certainly, the available studies indicate that plants like *Ageratum conyzoides*, present excellent potential for development of drugs leading to treatment of both widespread but common ailments like gastrointestinal disorders, as well as complicated ailments like diabetes and heart diseases, which are prevalent throughout the world population and which cannot be treated satisfactorily with modern allopathic medicine.

Ocimum gratissimum showed a distinguished antibacterial activity against

E. coli and this is line with the observations of earlier workers. This study however, showed the minimum inhibitory concentration of *O. gratissimum* to be rather high. With improved technology, better active principle could be extracted from the plant. This could be of value in the treatment of diarrhea due to *E. coli* in livestock. In the study by Adebayo-Tayo and Adegoke, *O. gratissimum* and *H. floribunda* exhibited the highest degree of activity and activity index. In their study, the extract of *O. graissimum* ("Nton") was reported to have had the highest mean inhibitory zone of 23 mm against *S. flexineri*. In the same study by Adebayo-Tayo and Adegoke, the MIC results of *S. flexineri* in 26.24 mg/ml of *O. gratissimum* excite them to analyze for its MBC which was 47.24 mg/ml. However, the extracts showed the highest degree of inhibition against *S. flexineri* at concentration of 80 mg/ml, followed by *S. anceps* and the least was observed from *K. senegalensis* while at 40 mg/ml, *O. gratissimum* had the highest inhibitory concentration against *S. flexineri*.

In this study, *Ocimum gratissimum* has proven antibacterial properties useful in controlling diarrhea. Farmers should be encouraged to plant it on their farms. It can be fed to weaners. It is of value in controlling post-weaning diarrhea. Cattle farmers do not always avoid modern veterinary drugs because of their higher prices or lack of accessibility; but because in their actual observations, they found out that the use of medicinal plants for treatment of certain cattle ailments can provide better efficacy. Modern research has also started to validate at least some

of the veterinary uses of medicinal plants.

The activities of the plants extracts also varied between solvents with the methanol extracts demonstrating the highest activity against all the test bacteria. This is consistent with other studies. In this study, the methanol extracts of the plant parts were more potent than the chloroform and water extracts, which showed no inhibitory activity. This is similar to the reports of Elmahmood and Amey but contrary to observations of Roy et al. It has been reported that different phytoconstituents have different degrees of solubility in different types of solvents depending on their polarity. In a traditional setting, water is the solvent largely used to prepare these concoctions. The higher activity demonstrated by organic solvents in this study is therefore an indication that less of the bioactive principles are extracted when water is used as a solvent. In conclusion, the development of bacterial resistance to presently available antibiotics has necessitated the search for new antibacterial agents. The 3 plants (*Ageratum conyzoides*, *Annona muricata* and *Ocimum gratissimum*) that showed some degree of inhibitions with ethanol extract may be indicative of the presence of antibiotics compounds in them. In line with Yagoub, this may help to discover new phytochemical classes of antibiotic substances that could serve as selective agents for infectious disease chemotherapy and control. More investigation on the subject especially the effect of these 3 plants (*Ageratum conyzoides* and *Ocimum gratissimum*) on other agents of diarrhea such as

Salmonella spp and *Yersinia* spp will be rewarding. This investigation might aid the possibility of the use of plants in drug development for human consumption possibly for the treatment of gastrointestinal, urinary tract and wound infection as well as typhoid fever. The noted medicinal plants with no antibacterial activity on *E. coli* but which have been empirically proven to have anti-diarrhoeic effects may be due to astringent properties possessed by extract of the plants. This act by preventing loss of electrolytes and water to the intestinal lumen. The extracts may show antibacterial activities against other agents of diarrhea that have not been tested in this investigation. There is the need to carry out further test on other agents of diarrhea in order to know the spectrum of microorganisms that are susceptible to the different fractions of *Ageratum conyzoides* and *Ocimum gratissimum*. According to Yagoub, the effect of these plants on more pathogenic organisms and toxicological investigation and purification however, needs to be carried out. Thus, in line with the assertions of Harun-or-Rashid et al., our findings present considerable potential for further scientific research on these plant species, which can lead to development of cheaper and more efficacious drugs. However, toxicological analysis of the crude extracts is recommended in order to assess their safety in the body of a patient when administered.

REFERENCES

Corrêa MP 1932. Dicionário das Plantas Úteis do Brasil, IBDF, Ministério da

Agricultura, Rio de Janeiro, 63 pp.

Ilory M, Sheteolu AO, Omonibgehin EA, Adeneye AA 1996. Antidiarrhoeal activities of *Ocimum gratissimum* (Lamiaceae). *J Diarrhoeal Dis Res* 14: 283-285.

Janssen AM, Scheffer JJ, Ntezurubanza L, Baerheim Svendsen A 1989. Antimicrobial activities of some *Ocimum* species grown in Rwanda. *J Ethnopharmacol* 26: 57-63.

Jedlickova Z, Motti O, Sery V 1992. Antibacterial properties of the Vietnamese cajeput oil and ocimum oil in combination with anticaterial agents. *J Hyg Epidemiol Microbiol Immunol* 36: 303-309.

Lima EO, Gompetz OF, Giesbrecht AM, Paulo MQ 1993. In vitro antifungal activity of essential oils obtained from officinal plants against dermatophytes. *Mycoses* 36: 333-336.

Moffat AC, Jackson JV, Moss MS, Widdop B 1986. *Clarke's Isolation and Identification of Drugs*, The Pharmaceutical Press, London, 609 pp.

Nwosu MO, Okafor JI 1995. Preliminary studies of the antifungal activities of some medical plants against *Basidiobolus* and some other pathogenic fungi. *Mycoses* 38: 191-195.

Obaseiki-Ebor EE, Odukoya K, Telikepally H, Mitscher LA, Shankel DM 1993. Antimutagenic activity of extracts of leaves of four common edible vegetable plants in Nigeria (West Africa). *Mutat Res* 302: 109-117.

Oliver B 1960. *Medicinal Plants in Nigeria*, Nigerian College of Arts, Science and

Technology, Nigeria, 42 pp.

Onajobi FD 1986. Smooth muscle contracting lipid soluble principles in chromatographic fractions of *Ocimum gratissimum*. *J Ethnopharmacol* 18: 3-11.

Ramanoelina AR, Terrom GP, Bianchini JP, Coulanges P 1987. Antibacterial action of essential oils extracted from Madagascar plants. *Arch Inst Pasteur Madagascar* 53: 217-226.

Sainsbury M, Sofowora EA 1971. Essential oil from the leaves and inflorescence of *Ocimum gratissimum*. *Phytochemistry* 10: 3309-3310.

Stahl E 1969. *Thin-layer Chromatography*, Springer Verlag, New York, 904 pp.